

Paul Baird; Jade Ventura

Four-dimensional Einstein metrics from biconformal deformations

Archivum Mathematicum, Vol. 57 (2021), No. 5, 255–283

Persistent URL: <http://dml.cz/dmlcz/149132>

Terms of use:

© Masaryk University, 2021

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

FOUR-DIMENSIONAL EINSTEIN METRICS FROM BICONFORMAL DEFORMATIONS

PAUL BAIRD AND JADE VENTURA

ABSTRACT. Biconformal deformations take place in the presence of a conformal foliation, deforming by different factors tangent to and orthogonal to the foliation. Four-manifolds endowed with a conformal foliation by surfaces present a natural context to put into effect this process. We develop the tools to calculate the transformation of the Ricci curvature under such deformations and apply our method to construct Einstein 4-manifolds. Examples of one particular family have ends which collapse asymptotically to \mathbb{R}^2 .

1. INTRODUCTION

A smooth Riemannian manifold (M, g) is said to be *Einstein* if its Ricci curvature satisfies $\text{Ric} = Ag$ for some constant A . D. Hilbert showed how Einstein metrics arise from the variational problem of extremizing scalar curvature [8]. The relation between scalar curvature and conformal transformations has been explored by analysts over the latter part of the last century. The Yamabe problem is to determine the existence of a metric of constant scalar curvature in a conformal class [14]. There have been important contributions by various authors and the problem was completely solved positively in the compact case by R. Schoen [10]; for a survey see the notes of Hebey [6].

Conformal transformations are not in general sufficiently discerning to find Einstein metrics. For example, although any manifold admits a Riemannian metric, on a compact manifold, there is a topological obstruction to the existence of an Einstein metric, known as the Hitchin-Thorpe inequality [2, 9, 12], whereas there always exist constant scalar curvature metrics. Biconformal deformations on the other hand, appear optimal to control the Ricci curvature.

A biconformal deformation of a Riemannian manifold (M, g) (see below) takes place in the presence of a conformal foliation. A foliation \mathcal{F} is *conformal* if Lie transport along the leaves of the normal space is conformal [13], specifically, if we

2020 *Mathematics Subject Classification*: primary 53C25; secondary 53C18, 53C12.

Key words and phrases: Einstein manifold, conformal foliation, semi-conformal map, biconformal deformation.

The second author acknowledges the support of the Philippine Commission on Higher Education and Campus France. The images were created with GeoGebra by Dr. Adrian Jannetta - <https://www.geogebra.org/m/W7dAdgqc>.

Received August 20, 2021. Editor J. Slovák.

DOI: 10.5817/AM2021-5-255

set $T\mathcal{F}$ to be the tangent space to the leaves and $N\mathcal{F}$ the normal space, there exists a mapping $a: T\mathcal{F} \rightarrow \mathbb{R}$, linear at each point, such that

$$(\mathcal{L}_U g)(X, Y) = a(U)g(X, Y) \quad (\forall U \in T\mathcal{F}, \forall X, Y \in N\mathcal{F}).$$

Conformal foliations are intimately related to semi-conformal mappings.

A mapping $\varphi: (M^m, g) \rightarrow (N^n, h)$ is *semi-conformal* if at each point where its derivative is non-zero, it is surjective and conformal (and so homothetic) on the complement of its kernel. Specifically, at each $x \in M$ where $d\varphi_x \neq 0$, the derivative is surjective and there exists a real number $\lambda(x) > 0$ such that

$$\varphi^* h(X, Y) = \lambda(x)^2 g(X, Y) \quad (\forall X, Y \in (\ker d\varphi_x)^\perp).$$

Extending λ to be zero at points x where $d\varphi_x = 0$, determines a continuous function $\lambda: M \rightarrow \mathbb{R}(\geq 0)$, smooth away from critical points, called the *dilation* of φ . In [1], it is shown that if $\varphi: (M^m, g) \rightarrow (N^n, h)$ is a semi-conformal submersion, then its fibres form a conformal foliation; conversely, if \mathcal{F} is a conformal foliation on (M^m, g) and $\psi: W \subset M \rightarrow \mathbb{R}^n \times \mathbb{R}^{m-n}$ is a local foliated chart, then there is a conformal metric on the leaf space N of $\mathcal{F}|_W$ with respect to which the natural projection $\varphi: W \rightarrow N$ is a semi-conformal submersion. The relation between a above and the dilation λ is given by $a = -2d \ln \lambda|_{\mathcal{V}}$, where $\mathcal{V} = T\mathcal{F} = \ker d\varphi$ [1].

Let $\varphi: (M^n, g) \rightarrow (N^n, h)$ be a semi-conformal submersion between Riemannian manifolds. Then the metric g decomposes into the sum $g = g^H + g^V$ of its horizontal and vertical components. A *biconformal deformation* of g is a metric

$$\tilde{g} = \frac{g^H}{\sigma^2} + \frac{g^V}{\rho^2},$$

where $\sigma, \rho: M \rightarrow \mathbb{R}$ are smooth positive functions. Note that the deformation is conformal if and only if $\sigma \equiv \rho$. We could equally define a biconformal deformation with respect to a conformal foliation. Such deformations preserve semi-conformality of φ .

The idea to use biconformal deformations to construct 4-dimensional Einstein metrics is founded on the possibility of obtaining a suitable expression for the Ricci curvature in terms of parameters of the semi-conformal map: its dilation, second fundamental form of its fibres, integrability form associated to the horizontal distribution and the almost complex structure J given by rotation through $\pi/2$ in the horizontal and vertical spaces. When the mapping is a harmonic morphism with 1-dimensional fibres, an elegant expression was exploited by L. Danielo to construct Einstein metrics in dimension 4 by biconformally deforming the metric with respect to a harmonic morphism to a 3-manifold, with the deformation restricted to preserve harmonicity [3, 4].

In this article we achieve a computation of the Ricci curvature associated to a semi-conformal submersion $\varphi: (M^4, g) \rightarrow (N^2, h)$ (see §3) and use it to construct Einstein metrics by biconformal deformation associated to orthogonal projection $\mathbb{R}^4 \rightarrow \mathbb{R}^2$. Amongst the examples produced are warped product solutions deriving from a 3-dimensional dynamical system (see §5.1) and a family of complete Einstein metrics of negative Ricci curvature with each member having an \mathbb{R}^2 -end (Theorem

5.2). The term *end* is used loosely here to refer to a component of the exterior of a family of exhaustive subsets (not compact) that collapses to \mathbb{R}^2 .

In §2, we calculate the connection coefficients associated to a semi-conformal submersion $\varphi: (M^4, g) \rightarrow (N^2, h)$. We exploit these formulae in §3 to deduce expressions for the Ricci curvature in terms of the geometric parameters associated to φ referred to above. In §4, we obtain expressions for how these quantities change under biconformal transformation. These are then applied in §5 to orthogonal projection $\mathbb{R}^4 \rightarrow \mathbb{R}^2$, to deduce partial differential equations for an Einstein metric in terms of the parameters σ and ρ . In general these are challenging to solve, but special cases yield interesting and possibly new 4-dimensional Einstein metrics.

2. CONNECTION COEFFICIENTS ASSOCIATED TO A SEMI-CONFORMAL SUBMERSION

Let $\varphi: (M^4, g) \rightarrow (N^2, h)$ be a semi-conformal submersion between oriented Riemannian manifolds with dilation $\lambda: M \rightarrow \mathbb{R}^+$. The coefficients of the Levi-Civita connection with respect to an adapted orthonormal frame field will be expressed in terms of the dilation, the mean-curvature of the fibres and an integrability form associated to the horizontal distribution.

Let $\{f_1, f_2\}$ be a positive orthonormal frame on N^2 . Then in general $\nabla f_1 = \rho_{12}f_2$ and $\nabla f_2 = \rho_{21}f_1$ where $\rho_{12} = -\rho_{21}$ is the associated Cartan 1-form. Since the notion of semi-conformal is conformally invariant and since any Riemannian surface is locally conformally equivalent to a domain of \mathbb{R}^2 with its standard metric, for the rest of this section, we suppose the frame $\{f_1, f_2\}$ parallel, so the connection form ρ_{12} vanishes. By a trick, we will later remove this assumption in our expression for the Ricci curvature.

Let $\{e_1, e_2, e_3, e_4\}$ be a positive orthonormal frame on M^4 such that $d\varphi(e_i) = \lambda f_i$ for $i = 1, 2$, and $e_3, e_4 \in V := \ker d\varphi$. We will use indices in the following way: $i, j, \dots \in \{1, 2\}$, $r, s, \dots \in \{3, 4\}$, $a, b, \dots \in \{1, 2, 3, 4\}$ and sum over repeated indices. At each $x \in M$, let $\mathcal{H}_x: T_x M \rightarrow H_x = V_x^\perp$ denote orthogonal projection onto the horizontal space. If we don't wish to be specific about the point x we will simply write \mathcal{H} . Similarly, \mathcal{V} denotes projection onto the vertical space.

Define complementary indices i', j', \dots by $i' = 2$ if $i = 1$ and $i' = 1$ if $i = 2$. Set J^H to be rotation by $+\pi/2$ in the horizontal space H , thus: $J^H(e_1) = e_2$ and $J^H(e_2) = -e_1$, equivalently $J^H(e_i) = (-1)^{i+1}e_{i'}$. Similarly, set J^V to be rotation by $+\pi/2$ in the vertical space V , thus: $J^V(e_3) = e_4$ and $J^V(e_4) = -e_3$. Then $J := (J^H, J^V)$ defines an almost Hermitian structure on (M, g) .

Definition 2.1. For a semi-conformal submersion as above, define the *integrability 1-form* $\zeta: TM \rightarrow \mathbb{R}$ by

$$\zeta(X) := g(\nabla_{e_1} e_2, \mathcal{V}(X)) = \frac{1}{2}g([e_1, e_2], \mathcal{V}(X)) \quad \forall X \in TM,$$

where \mathcal{V} is orthogonal projection onto $\ker d\varphi$ and the second equality follows from *Lemma 2.4(i)* below. Then, ζ is well-defined independently of the (positive) horizontal orthonormal frame $\{e_1, e_2\}$ and vanishes if and only if the horizontal distribution is integrable.

Definition 2.2. Let $\mathcal{S} = \varphi^{-1}(y)$ be a fibre of φ . Then for vector fields X, Y tangent to \mathcal{S} , we have

$$\nabla_X Y = \nabla_X^{\mathcal{S}} Y + B_X Y$$

where ∇ is the connection on M , $\nabla^{\mathcal{S}}$ the connection on \mathcal{S} , i.e. $\nabla_X^{\mathcal{S}} Y = \mathcal{V}\nabla_X Y$, and B is the *second fundamental form of \mathcal{S}* (symmetric by integrability of the vertical distribution). Then the *mean curvature of the fibre* $\mu := \frac{1}{2}\text{Tr } B = \frac{1}{2}\mathcal{H}(\nabla_{e_3} e_3 + \nabla_{e_4} e_4)$.

Extend B to all vectors by the formula $B_X Y := \mathcal{H}\nabla_{\mathcal{V}X}\mathcal{V}Y$. Then its adjoint is characterized by:

$$g(B_X Y, Z) = g(Y, B_X^* Z) \quad \Rightarrow \quad B_X^* Z = -\mathcal{V}\nabla_{\mathcal{V}X}\mathcal{H}Z.$$

Lemma 2.3 (Fundamental equation of a semi-conformal submersion [1]). *For a semi-conformal submersion $\varphi: (M^m, g) \rightarrow (N^n, h)$, the tension field $\tau_\varphi = \text{Tr}_g \nabla d\varphi$ is given by*

$$\tau_\varphi = -(n-2)d\varphi(\text{grad } \ln \lambda) - (m-n)d\varphi(\mu)$$

where μ is the mean-curvature of the fibres.

Recall that the connection forms ω_{ab} are defined by $\nabla e_a = \sum_b \omega_{ab} e_b$. In order to express the connection coefficients, we require only the form ω_{34} . The following lemma expresses the connection coefficients in terms of the above quantities.

Lemma 2.4.

- (i) $\nabla_{e_i} e_j = -e_j(\ln \lambda) e_i + \text{grad } \ln \lambda + (-1)^{i+1} \delta_{ij'} \zeta^{\#}$
- (ii) $\nabla_{e_i} e_r = -e_r(\ln \lambda) e_i - \zeta(e_r) J e_i + \omega_{34}(e_i) J e_r$
- (iii) $\nabla_{e_r} e_i = -\zeta(e_r) J e_i - B_{e_r}^* e_i$
- (iv) $\nabla_{e_r} e_s = B_{e_r} e_s + \omega_{34}(e_r) J e_s.$

Proof. (i) From Lemma 2.3,

$$\tau_\varphi = -2d\varphi(\mu).$$

But, recalling we sum over repeated indices, $\nabla d\varphi(e_r, e_r) = -d\varphi(\nabla_{e_r} e_r) = -2d\varphi(\mu)$, so that

$$\nabla d\varphi(e_i, e_i) = \tau_\varphi - \nabla d\varphi(e_r, e_r) = 0.$$

On the other hand,

$$\begin{aligned} \nabla d\varphi(e_i, e_i) &= (-d\varphi(\nabla_{e_i} e_i) + \nabla_{e_i}^{\varphi^{-1}} d\varphi(e_i)) \\ &= (-d\varphi(\nabla_{e_i} e_i) + e_i(\ln \lambda) d\varphi(e_i) + \lambda^2 \nabla_{f_i}^N f_i) \\ &= (-d\varphi(\nabla_{e_i} e_i) + e_i(\ln \lambda) d\varphi(e_i)) = (-d\varphi(\nabla_{e_i} e_i) + e_i(\ln \lambda) d\varphi(e_i)). \end{aligned}$$

The expression for the horizontal component of $\nabla_{e_i} e_j$ now follows when we note that $g(e_1, \nabla e_1) = 0$ etc.

For the vertical component, first note that

$$(1) \quad g([e_r, e_i], e_j) = e_r(\ln \lambda) g(e_i, e_j) \quad (\forall i, j, \in \{1, 2\} \quad \forall r \in \{3, 4\}),$$

since, on the one hand

$$\nabla d\varphi(e_i, e_r) = -d\varphi(\nabla_{e_i} e_r);$$

on the other hand, by the symmetry of the second fundamental form

$$\begin{aligned}\nabla d\varphi(e_i, e_r) &= \nabla d\varphi(e_r, e_i) = -d\varphi(\nabla_{e_r} e_i) + \nabla_{e_r}^{\varphi^{-1}} d\varphi(e_i) \\ &= -d\varphi(\nabla_{e_r} e_i) + e_r(\ln \lambda) d\varphi(e_i) \\ &\implies d\varphi(\nabla_{e_i} e_r) = d\varphi(\nabla_{e_r} e_i) - e_r(\ln \lambda) d\varphi(e_i).\end{aligned}$$

Equation (1) follows. But then

$$\begin{aligned}-g(\nabla_{e_i} e_j, e_r) &= g(e_j, \nabla_{e_i} e_r) = g(e_j, \nabla_{e_r} e_i) - e_r(\ln \lambda) g(e_j, e_i) \\ -g(\nabla_{e_j} e_i, e_r) &= g(e_i, \nabla_{e_j} e_r) = g(e_i, \nabla_{e_r} e_j) - e_r(\ln \lambda) g(e_i, e_j).\end{aligned}$$

Now add and use the fact that $0 = e_r(g(e_i, e_j)) = g(\nabla_{e_r} e_i, e_j) + g(e_i, \nabla_{e_r} e_j)$.

(ii) follows since

$$\begin{aligned}\mathcal{H}\nabla_{e_i} e_r &= g(\nabla_{e_i} e_r, e_j) e_j = -g(e_r, \nabla_{e_i} e_j) e_j \\ &= -e_r(\ln \lambda) e_i + (-1)^i \zeta(e_r) e_{i'} = -e_r(\ln \lambda) e_i - \zeta(e_r) J^{\mathcal{H}} e_i.\end{aligned}$$

(iii) follows from (1) and (ii).

(iv) is a consequence of the definitions. \square

Corollary 2.5.

- (i) $[e_i, e_j] = e_i(\ln \lambda) e_j - e_j(\ln \lambda) e_i + 2(-1)^{i+1} \delta_{ij'} \zeta^{\sharp}$
- (ii) $[e_r, e_i] = e_r(\ln \lambda) e_i - B_{e_r}^* e_i - \omega_{34}(e_i) J e_r$
- (iii) $\nabla_{e_i} e_i = \text{grad } \ln \lambda + \mathcal{V} \text{grad } \ln \lambda$
- (iv) $\nabla_{e_a} e_a = \text{grad } \ln \lambda + \mathcal{V} \text{grad } \ln \lambda + 2\mu + \omega_{34}(e_r) J e_r.$

3. THE RICCI CURVATURE

Let $\varphi: (M^4, g) \rightarrow (N^2, h)$ be a semi-conformal submersion between oriented Riemannian manifolds. Choose an orthonormal frame field $\{e_a\} = \{e_i; e_r\}$ adapted to the horizontal and vertical spaces. The Ricci curvature is determined by its components:

$$\text{Ric} = R_{ab} \theta_a \theta_b = R_{11} \theta_1^2 + 2R_{12} \theta_1 \theta_2 + \dots$$

where $\{\theta_a\}$ is the dual frame to $\{e_a\}$ and the product $\theta_a \theta_b = \theta_a \odot \theta_b = \frac{1}{2}(\theta_a \otimes \theta_b + \theta_b \otimes \theta_a)$ is the symmetric product of 1-forms. The coefficients R_{ab} are symmetric in their indices and $R_{ab} = \text{Ric}(e_a, e_b)$. In order to compute the Ricci curvature associated to a semi-conformal submersion, we will separately calculate the horizontal components R_{ij} , the mixed components R_{ri} and the vertical components R_{rs} .

Define the covariant tensor fields C and C^* by

$$\begin{aligned}C(X, Y) &:= g(B_{e_r} X, B_{e_r} Y) = g(\text{Tr}(B^* B)(X), Y) \\ C^*(X, Y) &:= g(B_{e_r}^* X, B_{e_r}^* Y) = g(\text{Tr}(B B^*)(X), Y).\end{aligned}$$

Note that C and C^* are well-defined independent of the frame, symmetric and that C vanishes on horizontal vectors and C^* on vertical vectors.

For a general covariant tensor field $T(X, Y, Z, \dots)$, define its divergence as derivation and contraction with respect to the *first* entry:

$$(\operatorname{div} T)(Y, Z, \dots) = (\nabla_{e_a} T)(e_a, Y, Z, \dots) = e_a(T(e_a, Y, Z, \dots)) - T(\nabla_{e_a} e_a, Y, Z, \dots) \\ - T(e_a, \nabla_{e_a} Y, Z, \dots) - T(e_a, Y, \nabla_{e_a} Z, \dots) - \dots$$

To the second fundamental form of the fibres B (a $(2, 1)$ tensor field), we associate two $(3, 0)$ -tensor fields. The first of these is $B_1: TM \times TM \times TM \rightarrow \mathbb{R}$ determined by

$$B_1(X, Y, Z) = g(X, \mathcal{H}\nabla_{\mathcal{V}Y}\mathcal{V}Z)$$

and the second

$$B_2(X, Y, Z) = g(\mathcal{H}\nabla_{\mathcal{V}X}\mathcal{V}Y, Z).$$

Note that B_1 and B_2 are identical up to ordering of their arguments, however, their divergences differ.

Our aim is to calculate the Ricci curvature in terms of parameters associated to φ . Being a tensorial object, it suffices to calculate Ric at a point x_0 where we can suppose the frame chosen such that $\mathcal{V}\nabla_{e_r}e_s = 0$, for all $r, s = 3, 4$. Such a frame can be constructed by first choosing a local *normal* frame $\{e_r\}$ for the fibre $\varphi^{-1}(\varphi(x_0))$ centered on x_0 (see [11], Vol. 2, Chapter 7) and then extending this to an orthonormal frame $\{e_a\}$ about x_0 in M . In particular, at x_0 , we have $\omega_{34}(e_r) = 0$ for $r = 3, 4$.

Lemma 3.1. *Acting on vertical vectors, the divergence of B_1 at x_0 is determined by*

$$(\operatorname{div} B_1)(e_r, e_s) = e_i(g(e_i, B_{e_r}e_s)) - 2\mu^b(B_{e_r}e_s) - d \ln \lambda(B_{e_r}e_s) \\ - g(e_t, \nabla_{e_i}e_r)g(e_i, B_{e_t}e_s) - g(e_t, \nabla_{e_i}e_s)g(e_i, B_{e_r}e_t)$$

(recalling, we sum over repeated indices).

Proof.

$$(\operatorname{div} B_1)(e_r, e_s) = (\nabla_{e_a} B_1)(e_a, e_r, e_s) \\ = e_i(B_1(e_i, e_r, e_s)) - B_1(\nabla_{e_a} e_a, e_r, e_s) - B_1(e_i, \nabla_{e_i} e_r, e_s) \\ - B_1(e_i, e_r, \nabla_{e_i} e_s).$$

From Corollary 2.5(iv), at x_0 , $\mathcal{H}\nabla_{e_a}e_a = 2\mu + \mathcal{H}\operatorname{grad} \ln \lambda$; also $\mathcal{V}\nabla_{e_i}e_r = g(e_t, \nabla_{e_i}e_r)e_r$ etc. and the formula follows. \square

Lemma 3.2. *Acting on a vertical and a horizontal vector, the divergence of B_2 at x_0 is given by*

$$(\operatorname{div} B_2)(e_r, e_i) = e_s(B_2(e_s, e_r, e_i)) - 2g(B_{\mathcal{V}\operatorname{grad} \ln \lambda}e_r, e_i) - \zeta(\nabla_{e_r}J^{\mathcal{H}}e_i).$$

Proof. Calculating at x_0 ,

$$(\operatorname{div} B_2)(e_r, e_i) = e_a(B_2(e_a, e_r, e_i)) - B_2(\mathcal{V}\nabla_{e_a}e_a, e_r, e_i) - B_2(e_s, \mathcal{V}\nabla_{e_s}e_r, e_i) \\ - B_2(e_s, e_r, \mathcal{H}\nabla_{e_s}e_i) \\ = e_s(B_2(e_s, e_r, e_i)) - B_2(\mathcal{V}\nabla_{e_j}e_j, e_r, e_i) - B_2(e_s, e_r, \mathcal{H}\nabla_{e_s}e_i).$$

On applying Corollary 2.5(iii) and Lemma 2.4(iii), this becomes

$$e_s(B_2(e_s, e_r, e_i)) - 2g(B_{\mathcal{V}\text{grad } \ln \lambda} e_r, e_i) + \zeta(e_s)g(\nabla_{e_r} e_s, J^{\mathcal{H}} e_i).$$

But the latter term equals $-\zeta(e_s)g(e_s, \nabla_{e_r} J^{\mathcal{H}} e_i)$ and the formula follows. \square

In what follows, we shall first establish the stated formulae for the case when N^2 is flat; in particular, we can suppose that $d\varphi(e_i) = \lambda f_i$ where $\{f_i\}$ is a parallel frame: $\nabla f_i = 0$ and apply the formulae of §2. We will then extend the formulae to the case when N^2 is an arbitrary Riemannian surface.

3.1. The horizontal components of the Ricci curvature.

First, we require the following lemma.

Lemma 3.3. *The horizontal sectional curvature $K^H := g(R(e_1, e_2)e_2, e_1)$ is given by*

$$K^H = \Delta \ln \lambda - \text{Tr } \mathcal{V} \nabla d \ln \lambda + \|\mathcal{V} \text{grad } \ln \lambda\|^2 - 3\|\zeta\|^2.$$

Proof. From Lemma 2.4(i) and Corollary 2.5(i),

$$\begin{aligned} K^H &= g(\nabla_{e_1} \nabla_{e_2} e_2 - \nabla_{e_2} \nabla_{e_1} e_2 - \nabla_{[e_1, e_2]} e_2, e_1) \\ &= g(\nabla_{e_1} (e_1(\ln \lambda) e_1 + \mathcal{V} \text{grad } \ln \lambda) + \nabla_{e_2} (e_2(\ln \lambda) e_1 - \zeta^\sharp, e_1) \\ &\quad - e_1(\ln \lambda) g(\nabla_{e_2} e_2, e_1) + e_2(\ln \lambda) g(\nabla_{e_1} e_2, e_1) - 2g(\nabla_{\zeta^\sharp} e_2, e_1)) \\ &= e_1(e_1(\ln \lambda)) + e_2(e_2(\ln \lambda)) - \|\mathcal{V} \text{grad } \ln \lambda\|^2 - \|\zeta\|^2 \\ &\quad - e_1(\ln \lambda)^2 - e_2(\ln \lambda)^2 - 2\zeta(e_r)g(\nabla_{e_r} e_2, e_1) \\ &= \Delta(\ln \lambda) - \text{Tr } \mathcal{V} \nabla d \ln \lambda + d \ln \lambda(\nabla_{e_i} e_i) - \|\mathcal{V} \text{grad } \ln \lambda\|^2 \\ &\quad - \|\mathcal{H} \text{grad } \ln \lambda\|^2 - 3\|\zeta\|^2, \end{aligned}$$

which, from Corollary 2.5(iii), gives the required formula. \square

Lemma 3.4. *The horizontal part of the Ricci curvature: $\text{Ric}|_{H \times H}$ is given by*

$$\text{Ric}|_{H \times H} = \{\lambda^2 K^N + \Delta \ln \lambda + 2d \ln \lambda(\mu) - 2\|\zeta\|^2\} g^H - C^* + \mathcal{L}_\mu g|_{H \times H},$$

where K^N denotes the Gaussian curvature of N .

Proof. The horizontal components $R_{ij} = \text{Ric}(e_i, e_j)$ are given by

$$R_{ij} = g(R(e_i, e_a)e_a, e_j) = K^H g(e_i, e_j) + g(R(e_i, e_r)e_r, e_j)$$

where K^H is given by Lemma 3.3 above.

We now calculate $g(R(e_i, e_r)e_r, e_j) = g(\nabla_{e_i} \nabla_{e_r} e_r - \nabla_{e_r} \nabla_{e_i} e_r - \nabla_{[e_i, e_r]} e_r, e_j)$. Then

$$\begin{aligned} g(\nabla_{e_i} \nabla_{e_r} e_r, e_j) &= g(\nabla_{e_i} (\mathcal{H} \nabla_{e_r} e_r + \mathcal{V} \nabla_{e_r} e_r), e_j) \\ &= 2g(\nabla_{e_i} \mu, e_j) - g(\mathcal{V} \nabla_{e_r} e_r, \nabla_{e_i} e_j) = 2g(\nabla_{e_i} \mu, e_j). \end{aligned}$$

From Lemma 2.4(ii) and (iii),

$$\begin{aligned}
-g(\nabla_{e_r} \nabla_{e_i} e_r, e_j) &= -g(\nabla_{e_r} (\mathcal{H} \nabla_{e_i} e_r + \mathcal{V} \nabla_{e_i} e_r), e_j) \\
&= g(\nabla_{e_r} (e_r (\ln \lambda) e_i + \zeta(e_r) J e_i), e_j) - g(\mathcal{V} \nabla_{e_i} e_r, \nabla_{e_r} e_j) \\
&= e_r (e_r (\ln \lambda)) g(e_i, e_j) + e_r (\ln \lambda) g(\nabla_{e_r} e_i, e_j) \\
&\quad + g(\nabla_{e_r} (\zeta(e_r) J e_i), e_j) - g(\mathcal{V} \nabla_{e_i} e_r, \nabla_{e_r} e_j) \\
&= (\text{Tr}_V \nabla \ln \lambda + 2d \ln \lambda(\mu)) g(e_i, e_j) \\
&\quad + e_r (\ln \lambda) g(\nabla_{e_r} e_i, e_j) + g(\nabla_{e_r} (\zeta(e_r) J e_i), e_j) \\
&\quad - g(\mathcal{V} \nabla_{e_i} e_r, \nabla_{e_r} e_j).
\end{aligned}$$

From Lemma 2.4,

$$\begin{aligned}
[e_i, e_r] &= g([e_i, e_r], e_k) e_k + g([e_i, e_r], e_s) e_s \\
&= -e_r (\ln \lambda) e_i + g(\nabla_{e_i} e_r - \nabla_{e_r} e_i, e_s) e_s
\end{aligned}$$

so that

$$\begin{aligned}
-g(\nabla_{[e_i, e_r]} e_r, e_j) &= e_r (\ln \lambda) g(\nabla_{e_i} e_r, e_j) - g(\nabla_{e_i} e_r - \nabla_{e_r} e_i, e_s) g(\nabla_{e_s} e_r, e_j) \\
&= e_r (\ln \lambda) g(-e_r (\ln \lambda) e_i - \zeta(e_r) J e_i, e_j) \\
&\quad - g(\nabla_{e_i} e_r, e_s) g(\nabla_{e_s} e_r, e_j) + g(\nabla_{e_r} e_i, e_s) g(\nabla_{e_s} e_r, e_j) \\
&= -\|\mathcal{V} \text{grad } \ln \lambda\|^2 g(e_i, e_j) - e_r (\ln \lambda) \zeta(e_r) g(J e_i, e_j) \\
&\quad - g(\nabla_{e_i} e_r, e_s) g(\nabla_{e_s} e_r, e_j) + g(\nabla_{e_r} e_i, e_s) g(\nabla_{e_s} e_r, e_j).
\end{aligned}$$

However, the Ricci tensor is symmetric in its arguments: $\text{Ric}(e_i, e_j) = \frac{1}{2}(\text{Ric}(e_i, e_j) + \text{Ric}(e_j, e_i))$. But then $g(\nabla_{e_i} \mu, e_j) + g(\nabla_{e_j} \mu, e_i) = \mathcal{L}_\mu g(e_i, e_j)$, $g(J e_i, e_j) + g(J e_j, e_i) = 0$ and

$$\zeta(e_r) (g(\nabla_{e_r} J e_i, e_j) + g(\nabla_{e_r} J e_j, e_i)) = -\zeta(e_r) (g(J e_i, \nabla_{e_r} e_j) + g(J e_j, \nabla_{e_r} e_i)) = \|\zeta\|^2.$$

Collecting terms now gives the required expression in the case of flat codomain. \square

3.2. The mixed components of the Ricci curvature.

Lemma 3.5. *For X a horizontal vector and U a vertical vector, one has*

$$\begin{aligned}
\text{Ric}(X, U) &= \nabla d \ln \lambda(X, U) - (d \ln \lambda)^2(X, U) - 2(d \ln \lambda \odot \zeta)(JX, U) \\
&\quad - (\nabla_{JX} \zeta)(U) - 2\zeta(\nabla_U JX) - \text{div } B_2(U, X) \\
&\quad - 2d \ln \lambda(B_U^* X) + 2(\nabla_U \mu^b)(X).
\end{aligned}$$

Proof. By tensoriality, it suffices to set $X = e_i$ and $U = e_r$. Then

$$\text{Ric}(e_i, e_r) = g(R(e_i, e_a) e_a, e_r) = g(R(e_i, e_j) e_j, e_r) + g(R(e_r, e_s) e_s, e_i).$$

First, we deal term by term with

$$g(R(e_i, e_j) e_j, e_r) = g(\nabla_{e_i} \nabla_{e_j} e_j - \nabla_{e_j} \nabla_{e_i} e_j - \nabla_{[e_i, e_j]} e_j, e_r).$$

From Corollary 2.5(iii) and Lemma 2.4(ii),

$$\begin{aligned} g(\nabla_{e_i} \nabla_{e_j} e_j, e_r) &= g(\nabla_{e_i} (2\text{grad } \ln \lambda - \mathcal{H}\text{grad } \ln \lambda), e_r) \\ &= 2\nabla \text{d} \ln \lambda(e_i, e_r) + g(\mathcal{H}\text{grad } \ln \lambda, \nabla_{e_i} e_r) \\ &= 2\nabla \text{d} \ln \lambda(e_i, e_r) - e_i(\ln \lambda) e_r(\ln \lambda) - \zeta(e_r)(Je_i)(\ln \lambda). \end{aligned}$$

Also, from Lemma 2.4(ii),

$$\begin{aligned} -g(\nabla_{e_j} \nabla_{e_i} e_j, e_r) &= -e_j(g(\nabla_{e_i} e_j, e_r) + g(\nabla_{e_i} e_j, \nabla_{e_j} e_r)) \\ &= e_j(g(e_j, \nabla_{e_i} e_r)) + g(\nabla_{e_i} e_j, \nabla_{e_j} e_r) \\ &= e_j(-e_r(\ln \lambda)\delta_{ij} - \zeta(e_r)g(e_j, Je_i)) + g(\nabla_{e_i} e_j, \nabla_{e_j} e_r) \\ &= -e_i(e_r(\ln \lambda)) - (Je_i)(\zeta(e_r)) + g(\nabla_{e_i} e_j, \nabla_{e_j} e_r) \\ &= -e_i(e_r(\ln \lambda)) - (\nabla_{Je_i} \zeta)(e_r) - \zeta(\nabla_{Je_i} e_r) + g(\nabla_{e_i} e_j, \nabla_{e_j} e_r), \end{aligned}$$

where, from Lemma 2.4,

$$\begin{aligned} g(\nabla_{e_i} e_j, \nabla_{e_j} e_r) &= g(\nabla_{e_i} e_j, e_k)g(e_k, \nabla_{e_j} e_r) + g(\nabla_{e_i} e_j, e_s)g(e_s, \nabla_{e_j} e_r) \\ &= e_k(\ln \lambda)\delta_{ij}g(e_k, \nabla_{e_j} e_r) - e_j(\ln \lambda)\delta_{ik}g(e_k, \nabla_{e_j} e_r) \\ &\quad + e_s(\ln \lambda)\delta_{ij}g(e_s, \nabla_{e_j} e_r) + (-1)^{i+1}\delta_{ij'}\zeta(e_s)g(e_s, \nabla_{e_j} e_r) \\ &= g(\mathcal{H}\text{grad } \ln \lambda, \nabla_{e_i} e_r) - g(e_i, \nabla_{e_j} e_r)e_j(\ln \lambda) \\ &\quad + g(\mathcal{V}\text{grad } \ln \lambda, \nabla_{e_i} e_r) + \zeta(e_s)g(e_s, \nabla_{Je_i} e_r) \\ &= -2\zeta(e_r)\text{d} \ln \lambda(Je_i) + \text{d} \ln \lambda(\mathcal{V}\nabla_{e_i} e_r) + \zeta(\nabla_{Je_i} e_r). \end{aligned}$$

From Corollary 2.5(i) and Lemma 2.4(ii),

$$\begin{aligned} -g(\nabla_{[e_i, e_j]} e_j, e_r) &= -e_i(\ln \lambda)\delta_{jk}g(\nabla_{e_k} e_j, e_r) + e_j(\ln \lambda)\delta_{ik}g(\nabla_{e_k} e_j, e_r) \\ &\quad + 2(-1)^i\delta_{ij'}\zeta(e_s)g(\nabla_{e_s} e_j, e_r) \\ &= -2e_i(\ln \lambda)e_r(\ln \lambda) - g(\mathcal{H}\text{grad } \ln \lambda, \nabla_{e_i} e_r) \\ &\quad - 2\zeta(e_s)g(\nabla_{e_s} Je_i, e_r) \\ &= -e_i(\ln \lambda)e_r(\ln \lambda) + \zeta(e_r)\text{d} \ln \lambda(Je_i) - 2\zeta(\nabla_{e_r} Je_i). \end{aligned}$$

Collecting terms now yields

$$\begin{aligned} g(R(e_i, e_j)e_j, e_r) &= \nabla \text{d} \ln \lambda(e_i, e_r) - e_i(\ln \lambda)e_r(\ln \lambda) - \zeta(e_r)\text{d} \ln \lambda(Je_i) \\ &\quad - (\nabla_{Je_i} \zeta)(e_r) - 2\zeta(\nabla_{e_r} Je_i). \end{aligned}$$

For the other term, first note that at the point x_0 ,

$$\begin{aligned} g(\nabla_{e_r} \nabla_{e_s} e_s, e_i) &= g(\nabla_{e_r} (\mathcal{H}\nabla_{e_s} e_s + \mathcal{V}\nabla_{e_s} e_s), e_i) = 2g(\nabla_{e_r} \mu, e_i) \\ &\quad - g(\mathcal{V}\nabla_{e_s} e_s, \nabla_{e_r} e_i) = 2g(\nabla_{e_r} \mu, e_i). \end{aligned}$$

Then from Lemma 3.2,

$$\begin{aligned}
 g(R(e_r, e_s)e_s, e_i) &= g(\nabla_{e_r}\nabla_{e_s}e_s - \nabla_{e_s}\nabla_{e_r}e_s - \nabla_{[e_r, e_s]}e_s, e_i) \\
 &= 2g(\nabla_{e_r}\mu, e_i) - e_s(g(\nabla_{e_r}e_s, e_i)) + g(\nabla_{e_r}e_s, \nabla_{e_s}e_i) \\
 &\quad - g(\nabla_{[e_r, e_s]}e_s, e_i) \\
 &= 2(\nabla_{e_r}\mu^\flat)(e_i) - (\operatorname{div} B_2)(e_r, e_i) - 2g(\nabla_{e_r}\mathcal{V}\operatorname{grad} \ln \lambda, e_i) \\
 &\quad - \zeta(\nabla_{e_r}Je_i) + g(\mathcal{H}\nabla_{e_r}e_s, \mathcal{H}\nabla_{e_r}e_i).
 \end{aligned}$$

But from Lemma 2.4, $g(\mathcal{H}\nabla_{e_r}e_s, \mathcal{H}\nabla_{e_r}e_i) = -g(\nabla_{e_r}e_s, \zeta(e_s)Je_i) = \zeta(\nabla_{e_r}Je_i)$. The formula now follows for flat codomain. \square

3.3. The vertical components of the Ricci curvature.

Define the vertical sectional curvature by $K^V := g(R^F(e_3, e_4)e_4, e_3)$ where $F = \varphi^{-1}(y) \subset M$ is the fibre over $y \in N$ and R^F is the Riemannian curvature of F . Then K^V is related to the sectional curvature in M via the Gauss equation (see [11] Chapter 7):

$$g(R(e_3, e_4)e_4, e_3) = g(R^F(e_3, e_4)e_4, e_3) + |B_{e_3}e_4|^2 - g(B_{e_3}e_3, B_{e_4}e_4).$$

The correction terms have an invariant expression given by the following lemma, established by evaluating the right-hand and left-hand sides on the various (e_r, e_s) .

Lemma 3.6.

$$(|B_{e_3}e_4|^2 - g(B_{e_3}e_3, B_{e_4}e_4))g^V = C - 2\mu^\flat(B_\star\star).$$

Lemma 3.7.

$$\operatorname{Ric}|_{V \times V} = K^V g^V + 2\nabla \operatorname{d} \ln \lambda|_{V \times V} + 2\operatorname{d} \ln \lambda(B_\star\star) - 2(\operatorname{d} \ln \lambda)^2|_{V \times V} + 2\zeta^2 + \operatorname{div} B_1|_{V \times V}.$$

Proof.

$$\begin{aligned}
 \operatorname{Ric}(e_r, e_s) &= g(R(e_r, e_a)e_a, e_s) = (K^V + |B_{e_3}e_4|^2 \\
 &\quad - g(B_{e_3}e_3, B_{e_4}e_4))g(e_r, e_s) + g(R(e_r, e_i)e_i, e_s),
 \end{aligned}$$

with

$$g(R(e_r, e_i)e_i, e_s) = g(\nabla_{e_r}\nabla_{e_i}e_i - \nabla_{e_i}\nabla_{e_r}e_i - \nabla_{[e_r, e_i]}e_i, e_s).$$

From Corollary 2.5(iii), $\nabla_{e_i}e_i = \operatorname{grad} \ln \lambda + \mathcal{V}\operatorname{grad} \ln \lambda = 2\operatorname{grad} \ln \lambda - \mathcal{H}\operatorname{grad} \ln \lambda$, so that

$$\begin{aligned}
 g(\nabla_{e_r}\nabla_{e_i}e_i, e_s) &= 2g(\nabla_{e_r}\operatorname{grad} \ln \lambda, e_s) + g(\mathcal{H}\operatorname{grad} \ln \lambda, \nabla_{e_r}e_s) \\
 &= 2\nabla \operatorname{d} \ln \lambda(e_r, e_s) + \operatorname{d} \ln \lambda(B_{e_r}e_s).
 \end{aligned}$$

From Lemma 3.1,

$$\begin{aligned}
 -g(\nabla_{e_i}\nabla_{e_r}e_i, e_s) &= -e_i(g(\nabla_{e_r}e_i, e_s) + g(\nabla_{e_r}e_i, \nabla_{e_i}e_s)) \\
 &= \operatorname{div} B_1(e_r, e_s) + 2\mu^\flat(B_{e_r}e_s) + \operatorname{d} \ln \lambda(B_{e_r}e_s) \\
 &\quad + g(\nabla_{e_r}e_i, e_j)g(e_j, \nabla_{e_i}e_s) + g(\nabla_{e_r}e_i, e_t)g(e_t, \nabla_{e_i}e_s) \\
 &= \operatorname{div} B_1(e_r, e_s) + 2\mu^\flat(B_{e_r}e_s) + \operatorname{d} \ln \lambda(B_{e_r}e_s) \\
 &\quad + g(e_t, \nabla_{e_i}e_r)g(e_i, \nabla_{e_t}e_s) + g(\nabla_{e_r}e_i, e_j)g(e_j, \nabla_{e_i}e_s),
 \end{aligned}$$

where the last term can be expressed using Lemma 2.4(ii) and (iii):

$$g(\nabla_{e_r} e_i, e_j)g(e_j, \nabla_{e_i} e_s) = 2\zeta(e_r)\zeta(e_s).$$

From Corollary 2.5(ii) and (iii)

$$\begin{aligned} -g(\nabla_{[e_r, e_i]} e_i, e_s) &= -2e_r(\ln \lambda)e_s(\ln \lambda) - g(e_i, B_{e_r} e_t)g(e_i, B_{e_t} e_s) \\ &\quad + g(e_t, \nabla_{e_i} e_r)g(\nabla_{e_t} e_i, e_s). \end{aligned}$$

On collecting terms and applying Lemma 3.6, the formula follows for the case of flat codomain. \square

3.4. Mapping into an arbitrary curved surface.

Suppose $\varphi: (M^4, g) \rightarrow (N^2, h)$ is a semi-conformal submersion into an arbitrary Riemannian surface with dilation λ . About a point in the image of φ , choose local isothermal coordinates $\psi: W \rightarrow \mathbb{R}^2$ on an open set $W \subset N^2$, so that $h = \nu^{-2}(dy_1^2 + dy_2^2)$ for some function $\nu: W \rightarrow \mathbb{R}$. Consider the following composition:

$$(M^4, g) \xrightarrow{\varphi} (W \subset N^2, h) \xrightarrow{\psi} (W' \subset \mathbb{R}^2, \bar{h})$$

where \bar{h} is the canonical metric $dy_1^2 + dy_2^2$ on \mathbb{R}^2 and $W' = \psi(W)$. Then the formulae of §3.1, §3.2 and §3.3 apply to $\psi \circ \varphi$. We now show how they extend to φ .

Lemma 3.8.

$$\lambda^2 K^N \circ \varphi = \Delta \ln(\nu \circ \varphi) + 2d \ln(\nu \circ \varphi)(\mu).$$

Proof. First note that $K^N = \nu^{-2} \Delta_{\bar{h}} \ln \nu = \Delta_h \ln \nu$. Then from Lemma 2.3,

$$\begin{aligned} \Delta_g(\ln \nu \circ \varphi) &= d \ln \nu(\tau_\varphi) + \text{Tr}_g \nabla d \ln \nu(d\varphi, d\varphi) \\ &= -2d(\ln \nu \circ \varphi)(\mu) + \lambda^2(\Delta_h \ln \nu) \circ \varphi \\ &= -2d(\ln \nu \circ \varphi)(\mu) + \lambda^2 K^N \circ \varphi. \end{aligned} \quad \square$$

Since the dilation of $\psi \circ \varphi$ is given by $\lambda\nu$, from Lemma 3.4 (for the flat case),

$$\text{Ric}|_{H \times H} = \left\{ \Delta \ln(\lambda\nu) + 2d \ln(\lambda\nu)(\mu) - 2\|\zeta\|^2 \right\} g^{\mathcal{H}} - C^* + \mathcal{L}_\mu g|_{H \times H}.$$

But from Lemma 3.8,

$$\Delta \ln(\lambda\nu) + 2d \ln(\lambda\nu)(\mu) = \lambda^2 K^N + \Delta \ln \lambda + 2d \ln \lambda(\mu),$$

where the latter quantity is invariant with respect to conformal changes of metric on the codomain.

For the mixed components of the Ricci curvature, we note that on setting $\bar{\lambda} = \lambda\nu$,

$$\begin{aligned} &\nabla d \ln \lambda(X, U) - (d \ln \lambda)^2(X, U) - 2(d \ln \lambda \odot \zeta)(JX, U) \\ &= \nabla d \ln \bar{\lambda}(X, U) - (d \ln \bar{\lambda})^2(X, U) - 2(d \ln \bar{\lambda} \odot \zeta)(JX, U). \end{aligned}$$

For example

$$\nabla d \ln \bar{\lambda}(e_1, e_3) = \nabla d \ln \lambda(e_1, e_3) - d \ln(\nu \circ \varphi)(\nabla_{e_1} e_3).$$

But from Lemma 2.4,

$$\begin{aligned} -d \ln(\nu \circ \varphi)(\nabla_{e_1} e_3) &= -d \ln(\nu \circ \varphi)(\mathcal{H} \nabla_{e_1} e_3) \\ &= d \ln(\nu \circ \varphi)(g(e_3, \nabla_{e_1} e_1) e_1 + g(e_3, \nabla_{e_1} e_2) e_2) \\ &= 2(d \ln \lambda \odot d \ln(\nu \circ \varphi))(e_1, e_3) + 2(d \ln(\nu \circ \varphi) \odot \zeta)(J e_1, e_3). \end{aligned}$$

Whereas

$$\begin{aligned} &-(d \ln \bar{\lambda})^2(e_1, e_3) - 2(d \ln \bar{\lambda} \odot \zeta)(J e_1, e_3) \\ &= -(d \ln \lambda)^2(e_1, e_3) - 2(d \ln \lambda \odot \zeta)(J e_1, e_3) \\ &\quad - 2(d \ln \lambda \odot d \ln(\alpha \circ \varphi))(e_1, e_3) - 2(d \ln(\alpha \circ \varphi) \odot \zeta)(J e_1, e_3). \end{aligned}$$

The invariance of the vertical components of the Ricci curvature follows from the invariance of the quantity $\nabla d \ln \lambda|_{\mathcal{V} \times \mathcal{V}} + d \ln \lambda(B_\star \star)$, specifically $\nabla d \ln \lambda(e_r, e_s) + d \ln \lambda(B_{e_r} e_s) = e_r(e_s(\ln \lambda)) - d \ln \lambda(\mathcal{V} \nabla_{e_r} e_s) = e_r(e_s(\ln \bar{\lambda})) - d \ln \bar{\lambda}(\mathcal{V} \nabla_{e_r} e_s)$.

4. BICONFORMAL DEFORMATIONS

4.1. The effect of a biconformal deformation on the Ricci curvature.

Let $\varphi: (M^4, g_0) \rightarrow (N^2, h)$ be a semi-conformal map between oriented manifolds. Consider a biconformal deformation:

$$g = \frac{g_0^H}{\sigma^2} + \frac{g_0^V}{\rho^2}$$

where $\sigma, \rho: M^4 \rightarrow \mathbb{R}$ are smooth strictly positive functions. Write objects with respect to g_0 with an index 0, either upstairs or downstairs, and objects with respect to g as before. For example, the positive orthonormal basis with respect to g_0 will be written $\{e_1^0, e_2^0, e_3^0, e_4^0\}$ and the dilation of φ with respect to g_0 as λ_0 , etc. Then the new frame field and the dual field of 1-forms are given by

$$\begin{aligned} e_1 &= \sigma e_1^0, \quad e_2 = \sigma e_2^0, \quad e_3 = \rho e_3^0, \quad e_4 = \rho e_4^0 \\ \theta_1 &= \frac{1}{\sigma} \theta_1^0, \quad \theta_2 = \frac{1}{\sigma} \theta_2^0, \quad \theta_3 = \frac{1}{\rho} \theta_3^0, \quad \theta_4 = \frac{1}{\rho} \theta_4^0. \end{aligned}$$

The following lemma gives the change in the connection coefficients.

- Lemma 4.1.**
- (i) $g(\nabla_{e_r} e_s, e_i) = g_0(\nabla_{e_r}^0 e_s^0, e_i) + e_i(\ln \rho) \delta_{rs}$
 - (ii) $g(\nabla_{e_i} e_r, e_s) = g_0(\nabla_{e_i}^0 e_r^0, e_s^0)$
 - (iii) $g(\nabla_{e_r} e_i, e_j) = g_0(\nabla_{e_r}^0 e_i^0, e_j^0) + \frac{\rho^2 - \sigma^2}{2\rho^2} g_0([e_i^0, e_j^0], e_r)$
 - (iv) $g(\nabla_{e_r} e_s, e_t) = g_0(\nabla_{e_r}^0 e_s^0, e_t) + e_t(\ln \rho) \delta_{rs} - e_s(\ln \rho) \delta_{rt}$
 - (v) $g(e_i, \nabla_{e_k} e_j) = \sigma g_0(e_i^0, \nabla_{e_k}^0 e_j^0) + \sigma(e_i^0(\ln \sigma) \delta_{jk} - e_j^0(\ln \sigma) \delta_{ik})$
 - (vi) $g(\nabla_{e_i} e_j, e_r) = \frac{\sigma^2}{\rho^2} g_0(\nabla_{e_i}^0 e_j^0, e_r) + \left(1 - \frac{\sigma^2}{\rho^2}\right) e_r(\ln \lambda_0) \delta_{ij} + e_r(\ln \sigma) \delta_{ij}.$

Proof.

- (i) $2g(\nabla_{e_r} e_s, e_i) = g([e_i, e_r], e_s) + g([e_i, e_s], e_r)$
 $= \sigma g_0([e_i^0, e_r^0], e_s^0) + \sigma g_0([e_i^0, e_s^0], e_r^0) + 2\sigma e_i^0(\ln \rho) \delta_{rs}$
 $= 2\sigma g_0(\nabla_{e_r}^0 e_s^0, e_i^0) + 2\sigma e_i^0(\ln \rho) \delta_{rs}$
- (ii) $2g(\nabla_{e_i} e_r, e_s) = g([e_i, e_r], e_s) - g([e_i, e_s], e_r)$
 $= \frac{1}{\rho^2} g_0([e_i, \rho e_r^0], \rho e_s^0) - \frac{1}{\rho^2} g_0([e_i, \rho e_s^0], \rho e_r^0)$
 $= g_0([e_i, e_r^0], e_s^0) - g_0([e_i, e_s^0], e_r^0) + e_i(\ln \rho) \delta_{rs} - e_i(\ln \rho) \delta_{rs}$
 $= 2g_0(\nabla_{e_i} e_r^0, e_s^0)$
- (iii) $2g(\nabla_{e_r} e_i, e_j) = g([e_r, e_i], e_j) - g([e_i, e_j], e_r) + g([e_j, e_r], e_i)$
 $= \frac{1}{\sigma} g_0([\rho e_r^0, e_i^0], e_j^0) - \frac{1}{\rho} g_0([\sigma e_i^0, \sigma e_j^0], e_r^0) + \frac{1}{\sigma} g_0([\sigma e_j^0, \rho e_r^0], e_i^0)$
 $= \rho g_0([e_r^0, e_i^0], e_j^0) + \frac{\rho}{\sigma} e_r^0(\sigma) \delta_{ij} - \frac{\sigma^2}{\rho} g_0([e_i^0, e_j^0], e_r^0)$
 $+ \rho g_0([e_j^0, e_r^0], e_i^0) - \frac{\rho}{\sigma} e_r^0(\sigma) \delta_{ij}$
 $= 2\rho g_0(\nabla_{e_r}^0 e_i^0, e_j^0) + \frac{\rho^2 - \sigma^2}{\rho} g_0([e_i^0, e_j^0], e_r^0)$
- (iv) As above, we write $2g(\nabla_{e_r} e_s, e_t) = g([e_r, e_s], e_t) - g([e_s, e_t], e_r) + g([e_t, e_r], e_s)$ and replace e_r by ρe_r^0 etc. Case (v) is similar.
- (vi) $2g(\nabla_{e_i} e_j, e_r) = g([e_i, e_j], e_r) - g([e_j, e_r], e_i) + g([e_r, e_i], e_j)$
 $= \frac{1}{\rho} g_0([\sigma e_i^0, \sigma e_j^0], e_r^0) - \frac{1}{\sigma} g_0([\sigma e_j^0, \rho e_r^0], e_i^0) + \frac{1}{\sigma} g_0([\rho e_r^0, \sigma e_i^0], e_j^0)$
 $= 2\frac{\sigma^2}{\rho} g_0(\nabla_{e_i}^0 e_j^0, e_r^0) + \frac{\sigma^2}{\rho} (g_0([e_j^0, e_r^0], e_i^0) - g_0([e_r^0, e_i^0], e_j^0))$
 $- \rho g_0([e_j^0, e_r^0], e_i^0) + \rho g_0([e_r^0, e_i^0], e_j^0) + \rho e_r^0(\ln \sigma) \delta_{ij}$
 $+ \rho e_r^0(\ln \sigma) \delta_{ij}$

From Lemma 2.4, this gives

$$2g(\nabla_{e_i} e_j, e_r) = 2\frac{\sigma^2}{\rho} g_0(\nabla_{e_i}^0 e_j^0, e_r^0) - 2\frac{\sigma^2}{\rho} e_r^0(\ln \lambda_0) \delta_{ij} + 2\rho e_r^0(\ln \lambda_0) \delta_{ij} + 2\rho e_r^0(\ln \sigma) \delta_{ij}$$

and the formula follows. \square

Corollary 4.2.

- (i) $\nabla_{e_s} e_j = \sigma \rho \nabla_{e_r}^0 e_j^0 + \frac{\rho^2 - \sigma^2}{2\rho^2} \zeta^0(e_s) J e_j - e_j(\ln \rho) e_s$
- (ii) $\nabla_{e_r} e_s = \sigma^2 \mathcal{H} \nabla_{e_r}^0 e_s^0 + \rho^2 \mathcal{V} \nabla_{e_r}^0 e_s^0 + \delta_{rs} (\sigma^2 \mathcal{H} \text{grad}_{g_0} \ln \rho + \rho^2 \mathcal{V} \text{grad}_{g_0} \ln \rho)$
 $- \rho^2 e_s^0(\ln \rho) e_r^0.$

Proof. From Lemma 4.1,

$$\begin{aligned} \nabla_{e_s} e_j &= g(\nabla_{e_s} e_j, e_i) e_i + g(\nabla_{e_s} e_j, e_r) e_r = g(\nabla_{e_s} e_j, e_i) e_i - g(e_j, \nabla_{e_s} e_r) e_r \\ &= g_0(\nabla_{e_s}^0 e_j^0, e_i^0) e_i + \frac{\rho^2 - \sigma^2}{2\rho^2} g_0([e_j^0, e_i^0], e_s) e_i - g_0(\nabla_{e_s}^0 e_r^0, e_j) e_r - e_j(\ln \rho) \delta_{rs} e_r \\ &= \sigma \rho \nabla_{e_s}^0 e_j^0 + \frac{\rho^2 - \sigma^2}{2\rho^2} \zeta^0(e_s) J e_j - e_j(\ln \rho) e_s. \end{aligned}$$

The proof of (ii) is similar. \square

Corollary 4.3.

$$B_{e_r} e_s = \sigma^2 (B_{e_r}^0 e_s^0 + g_0(e_r^0, e_s^0) \mathcal{H} \text{grad}_{g_0} \ln \rho).$$

Proof. From (i) of Lemma 4.1,

$$g_0(B_{e_r}e_s, e_i^0) = \sigma^2(g_0(\nabla_{e_r}^0 e_s^0, e_i^0) + e_i^0(\ln \rho)\delta_{rs})$$

from which the formula follows. \square

Lemma 4.4. *The mean curvature of the fibres, the integrability form and the dilation change according to*

$$\mu = \sigma^2(\mu_0 + \mathcal{H}\text{grad}_{g_0} \ln \rho), \quad \zeta = \frac{\sigma^2}{\rho^2}\zeta_0, \quad \lambda = \sigma\lambda_0.$$

Proof. The expression for μ follows by taking the trace in Corollary 4.3. The Lie bracket is defined independently of the metric and the change in ζ follows. The expression for λ follows since the new horizontal basis is a multiple of σ times the old. \square

Lemma 4.5. *For a smooth function f ,*

$$\text{grad}_g f = \sigma^2 \text{grad}_{g_0} f + (\rho^2 - \sigma^2) \mathcal{V} \text{grad}_{g_0} f.$$

Proof.

$$\text{grad}_g f = e_a(f)e_a = \sigma^2 e_i^0(f)e_i^0 + \rho^2 e_r^0(f)e_r^0 = \sigma^2 \text{grad}_{g_0} f + (\rho^2 - \sigma^2) \mathcal{V} \text{grad}_{g_0} f.$$

\square

Recall that the basis $\{e_a^0\}$ is chosen such that at the point x_0 , we have $\mathcal{V}\nabla_{e_r}^0 e_s^0 = 0$, $\forall r, s = 3, 4$.

Lemma 4.6. *At the point x_0 ,*

$$\nabla_{e_a} e_a = \sigma^2(2\mu_0 + \mathcal{H}\text{grad}_{g_0} \ln(\sigma\lambda_0\rho^2)) + \rho^2 \mathcal{V} \text{grad}_{g_0} \ln(\rho\sigma^2\lambda_0^2)$$

Proof. From Corollary 2.5(iv),

$$\nabla_{e_a} e_a = \text{grad} \ln \lambda + \mathcal{V} \text{grad} \ln \lambda + 2\mu + \omega_{34}(e_r)Je_r.$$

From Lemma 4.1, $\omega_{34}(e_r)Je_r = \mathcal{V} \text{grad} \ln \rho$. The formula now follows from Lemmas 4.4 and 4.5. \square

Define the vertical Laplacian at a point x with respect to the metric g of a smooth function f by $\Delta_g^V f = \Delta_g^F(f|_F) = e_r(e_r(f)) - \text{d}f(\mathcal{V}\nabla_{e_r} e_r)$, where $F = \varphi^{-1}\varphi(x)$ is the fibre passing through x . Similarly, we have the vertical Laplacian with respect to g_0 . Note that at the point x_0 , we have $\Delta_{g_0}^V f = e_r^0(e_r^0(f))$.

Lemma 4.7. [3, 4]

$$\begin{aligned} \Delta_g f &= \sigma^2 \Delta_{g_0} f + (\rho^2 - \sigma^2) \{ \Delta_{g_0}^V f - 2\text{d}f(\mathcal{V} \text{grad}_{g_0} \ln \lambda_0) \} \\ &\quad - 2\sigma^2 \text{d}f(\mathcal{H} \text{grad}_{g_0} \ln \rho) - 2\rho^2 \text{d}f(\mathcal{V} \text{grad}_{g_0} \ln \sigma). \end{aligned}$$

Remark 4.8. Note that if $\sigma = \rho$, so that the transformation is conformal, we obtain the well-known formula for the transformation of the Laplacian:

$$\Delta_g f = \sigma^2 \Delta_{g_0} f - (m - 2)\sigma^2 \text{d}f(\text{grad}_{g_0} \ln \sigma)$$

(with dimension $m = 4$).

Proof. From Lemma 4.6,

$$\begin{aligned}
\Delta_g f &= e_a(e_a(f)) - \mathrm{d}f(\nabla_{e_a} e_a) = e_i(e_i(f)) + e_r(e_r(f)) \\
&\quad - \mathrm{d}f(2\sigma^2\mu_0 + \sigma^2\mathcal{H}\mathrm{grad}_{g_0} \ln \sigma \lambda_0 \rho^2 + \rho^2\mathcal{V}\mathrm{grad}_{g_0} \ln \rho \sigma^2 \lambda_0^2) \\
&= \sigma^2 e_i^0(e_i^0(f)) + \sigma^2 e_i^0(\ln \sigma) e_i^0(f) + \rho^2 e_r^0(e_r^0(f)) + \rho^2 e_r^0(\ln \rho) e_r^0(f) \\
&\quad - \mathrm{d}f(2\sigma^2\mu_0 + \sigma^2\mathcal{H}\mathrm{grad}_{g_0} \ln(\sigma \lambda_0 \rho^2) + \rho^2\mathcal{V}\mathrm{grad}_{g_0} \ln \rho \sigma^2 \lambda_0^2) \\
&= \sigma^2 \Delta_{g_0} f + (\rho^2 - \sigma^2) e_r^0(e_r^0(f)) + \sigma^2 \mathrm{d}f(\nabla_{e_a}^0 e_a^0) + \sigma^2 \mathrm{d}f(\mathcal{H}\mathrm{grad}_{g_0} \ln \sigma) \\
&\quad + \rho^2 \mathrm{d}f(\mathcal{V}\mathrm{grad}_{g_0} \ln \rho) \\
&\quad - 2\sigma^2 \mathrm{d}f(\mu_0) - \sigma^2 \mathrm{d}f(\mathcal{H}\mathrm{grad}_{g_0} \ln(\sigma \lambda_0 \rho^2)) - \rho^2 \mathrm{d}f(\mathcal{V}\mathrm{grad}_{g_0} \ln \rho \sigma^2 \lambda_0^2).
\end{aligned}$$

But from Corollary 2.5(iv), $\nabla_{e_a}^0 e_a^0 = \mathrm{grad}_{g_0} \ln \lambda_0 + \mathcal{V}\mathrm{grad}_{g_0} \ln \lambda_0 + 2\mu_0$, so that

$$\begin{aligned}
\Delta_g f &= \sigma^2 \Delta_{g_0} f + (\rho^2 - \sigma^2) e_r^0(e_r^0(f)) + \sigma^2 \mathrm{d}f(\mathrm{grad}_{g_0} \ln \lambda_0) + \sigma^2 \mathrm{d}f(\mathcal{V}\mathrm{grad}_{g_0} \ln \lambda_0) \\
&\quad + \sigma^2 \mathrm{d}f(\mathcal{H}\mathrm{grad}_{g_0} \ln \sigma) + \rho^2 \mathrm{d}f(\mathcal{V}\mathrm{grad}_{g_0} \ln \rho) - \sigma^2 \mathrm{d}f(\mathcal{H}\mathrm{grad}_{g_0} \ln(\sigma \lambda_0 \rho^2)) \\
&\quad - \rho^2 \mathrm{d}f(\mathcal{V}\mathrm{grad}_{g_0} \ln \rho \sigma^2 \lambda_0^2) \\
&= \sigma^2 \Delta_{g_0} f + (\rho^2 - \sigma^2) \{ \Delta_{g_0}^{\mathcal{V}} f - 2\mathrm{d}f(\mathcal{V}\mathrm{grad}_{g_0} \ln \lambda_0) \} \\
&\quad - 2\sigma^2 \mathrm{d}f(\mathcal{H}\mathrm{grad}_{g_0} \ln \rho) - 2\rho^2 \mathrm{d}f(\mathcal{V}\mathrm{grad}_{g_0} \ln \sigma).
\end{aligned}$$

□

Corollary 4.9.

$$\begin{aligned}
\Delta_g \ln \lambda &= \sigma^2 \Delta_{g_0} \ln(\sigma \lambda_0) + (\rho^2 - \sigma^2) \{ \Delta_{g_0}^{\mathcal{V}} (\ln(\sigma \lambda_0)) - 2\mathrm{d} \ln(\sigma \lambda_0) (\mathcal{V}\mathrm{grad}_{g_0} \ln \lambda_0) \} \\
&\quad - 2\sigma^2 \mathrm{d} \ln(\sigma \lambda_0) (\mathcal{H}\mathrm{grad}_{g_0} \ln \rho) - 2\rho^2 \mathrm{d} \ln(\sigma \lambda_0) (\mathcal{V}\mathrm{grad}_{g_0} \ln \sigma).
\end{aligned}$$

4.2. The second fundamental forms and their divergences. The vertical components of the Ricci tensor contain the term $\mathrm{div} B_1$ acting on vertical vectors.

Lemma 4.10.

$$B_1(e_i, e_r, e_s) = B_1^0(e_i, e_r^0, e_s^0) + \delta_{rs} e_i(\ln \rho).$$

Proof. This follows from Corollary 4.3:

$$\begin{aligned}
B_1(e_i, e_r, e_s) &= g(e_i, B_{e_r} e_s) = \frac{1}{\sigma^2} g_0(e_i, B_{e_r} e_s) \\
&= g_0(e_i, B_{e_r^0}^0 e_s^0 + g_0(e_r^0, e_s^0) \mathcal{H}\mathrm{grad}_{g_0} \ln \rho) \\
&= B_1^0(e_i, e_r^0, e_s^0) + \delta_{rs} e_i(\ln \rho).
\end{aligned}$$

□

Lemma 4.11.

$$\begin{aligned}
(\mathrm{div} B_1)(e_r, e_s) &= \sigma^2 (\mathrm{div}_0 B_1^0)(e_r^0, e_s^0) - \sigma^2 B_1^0(\mathcal{H}\mathrm{grad}_{g_0} \ln \rho^2, e_r^0, e_s^0) \\
&\quad + \delta_{rs} \sigma^2 \{ \mathrm{Tr}_{g_0}^H \nabla^0 \mathrm{d} \ln \rho + 2\mathrm{d} \ln \rho (\mathcal{V}\mathrm{grad}_{g_0} \ln \lambda_0) \\
&\quad - \mathrm{d} \ln \rho (2\mu_0 + \mathcal{H}\mathrm{grad}_{g_0} \ln \rho^2) \}.
\end{aligned}$$

(Note that $\mathrm{Tr}_{g_0}^H \nabla \mathrm{d} \ln \rho$ can be written in terms of the Laplacian and the vertical Laplacian).

Proof. Applying Lemma 4.10 and Lemma 4.6,

$$\begin{aligned}
(\operatorname{div} B_1)(e_r, e_s) &= (\nabla_{e_a} B_1)(e_a, e_r, e_s) = e_i(B_1(e_i, e_r, e_s)) - B_1(\nabla_{e_a} e_a, e_r, e_s) \\
&\quad - B_1(e_i, \nabla_{e_i} e_r, e_s) - B_1(e_i, e_r, \nabla_{e_i} e_s) \\
&= \sigma e_i^0(\sigma B_1^0(e_i^0, e_r^0, e_s^0) + \sigma \delta_{rs} e_i^0(\ln \rho)) \\
&\quad - \sigma^2 B_1(2\mu_0 + \mathcal{H}\operatorname{grad}_{g_0} \ln(\sigma \lambda_0 \rho^2), e_r, e_s) \\
&\quad - B_1(e_i, \nabla_{e_i} e_r, e_s) - B_1(e_i, e_r, \nabla_{e_i} e_s) \\
&= \sigma^2 e_i^0(B_1^0(e_i^0, e_r^0, e_s^0)) + \sigma^2 e_i^0(\ln \sigma) B_1^0(e_i^0, e_r^0, e_s^0) \\
&\quad + \delta_{rs} \sigma^2 e_i^0(e_i^0(\ln \rho)) + \delta_{rs} \sigma^2 e_i^0(\ln \sigma) e_i^0(\ln \rho) \\
&\quad - \sigma^2 B_1(2\mu_0, \mathcal{H}\operatorname{grad}_{g_0} \ln(\sigma \lambda_0 \rho^2), e_r, e_s) \\
&\quad - B_1(e_i, g_0(\nabla_{e_i}^0 e_r^0, e_t^0) e_t, e_s) - B_1(e_i, e_r, g_0(\nabla_{e_i}^0 e_s^0, e_t^0) e_t) \\
&= \sigma^2 \{ \operatorname{div}_0 B_1^0(e_r^0, e_s^0) + B_1^0(\nabla_{e_a^0}^0 e_a^0, e_r^0, e_s^0) + B_1^0(e_i^0, \nabla_{e_0^0}^0 e_r^0, e_s^0) \\
&\quad + B_1^0(e_i^0, e_r^0, \nabla_{e_0^0}^0 e_s^0) + B_1^0(\mathcal{H}\operatorname{grad}_{g_0} \ln \sigma, e_r^0, e_s^0) \\
&\quad + \delta_{rs} e_i^0(e_i^0(\ln \rho)) + \delta_{rs} e_i^0(\ln \sigma) e_i^0(\ln \rho) \\
&\quad - B_1^0(2\mu_0 + \mathcal{H}\operatorname{grad}_{g_0} \ln(\sigma \lambda_0 \rho^2), e_r^0, e_s^0) \\
&\quad - \delta_{rs} (2\mu_0 + \mathcal{H}\operatorname{grad}_{g_0} \ln(\sigma \lambda_0 \rho^2))(\ln \rho) \\
&\quad - B_1^0(e_i^0, \nabla_{e_0^0}^0 e_r^0, e_s^0) - B_1^0(e_i^0, e_r^0, \nabla_{e_0^0}^0 e_s^0) - (g_0(\nabla_{e_0^0}^0 e_s^0, e_r^0) \\
&\quad + g_0(\nabla_{e_0^0}^0 e_r^0, e_s^0)) e_i^0(\ln \rho) \}.
\end{aligned}$$

After simplifying and noting that from Corollary 2.5(iii)

$$\begin{aligned}
\operatorname{Tr}_{g_0}^H \nabla^0 \operatorname{div} \ln \rho &= -\operatorname{div} \ln \rho (\nabla_{e_0^0}^0 e_i^0) + e_i^0(e_i^0(\ln \rho)) \\
&= -\operatorname{div} \ln \rho (\operatorname{grad}_{g_0} \ln \lambda_0 + \mathcal{V}\operatorname{grad}_{g_0} \ln \lambda_0) + e_i^0(e_i^0(\ln \rho)),
\end{aligned}$$

the formula follows. \square

Let us now deal with $\operatorname{div} B_2$. As for Lemma 4.10, we have

Lemma 4.12.

$$B_2(e_r, e_s, e_i) = B_2^0(e_r^0, e_s^0, e_i) + \delta_{rs} e_i(\ln \rho).$$

Lemma 4.13.

$$\begin{aligned}
(\operatorname{div} B_2)(e_s, e_j) &= \operatorname{div}_{g_0} B_2^0(e_s, e_j) + \nabla^0 \operatorname{div} \ln \rho(e_s, e_j) - B_2^0(\mathcal{V}\operatorname{grad}_{g_0} \ln(\rho^3 \sigma), e_s, e_j) \\
&\quad - e_s(\ln(\sigma \lambda_0^2)) e_j(\ln \rho) + 2e_s(\ln \rho) \mu_0^b(e_j) \\
&\quad + \left(\frac{\sigma^2}{\rho^2} - 1 \right) B_2^0(\zeta_0^\sharp, e_s, J e_j) + \left(\frac{\sigma^2}{\rho^2} - 1 \right) \zeta_0(e_s) J e_j(\ln \rho).
\end{aligned}$$

Proof.

$$\begin{aligned}
(\operatorname{div} B_2)(e_s, e_j) &= (\nabla_{e_a} B_2)(e_a, e_s, e_j) = e_r(B_2(e_r, e_s, e_j)) \\
&\quad - B_2(\mathcal{V}\nabla_{e_a} e_a, e_s, e_j) - B_2(e_r, \mathcal{V}\nabla_{e_r} e_s, e_j) - B_2(e_r, e_s, \mathcal{H}\nabla_{e_r} e_j).
\end{aligned}$$

From Lemma 4.6, $\mathcal{V}\nabla_{e_a}e_a = \rho^2\mathcal{V}\text{grad}_{g_0}\ln\rho\sigma^2\lambda_0^2$. From Lemma 4.1(iv)

$$\mathcal{V}\nabla_{e_r}e_s = g(\nabla_{e_r}e_s, e_t)e_t = (e_t(\ln\rho)\delta_{rs} - e_s(\ln\rho)\delta_{rt})e_t = \delta_{rs}\mathcal{V}\text{grad}\ln\rho - e_s(\ln\rho)e_r,$$

and from Lemma 2.4(iii), $\mathcal{H}\nabla_{e_r}e_j = -\zeta(e_r)Je_j$. Thus, from Lemma 4.12,

$$\begin{aligned} (\text{div } B_2)(e_s, e_j) &= e_r(B_2(e_r, e_s, e_j)) - B_2(\rho^2\mathcal{V}\text{grad}_{g_0}\ln\rho\sigma^2\lambda_0^2, e_s, e_j) \\ &\quad - B_2(e_r, \delta_{rs}\mathcal{V}\text{grad}\ln\rho - e_s(\ln\rho)e_r, e_j) + \zeta(e_r)B_2(e_r, e_s, Je_j) \\ &= e_r(B_2(e_r, e_s, e_j)) - B_2(\rho^2\mathcal{V}\text{grad}_{g_0}\ln\rho\sigma^2\lambda_0^2, e_s, e_j) \\ &\quad + 2\mu^\flat(e_j)e_s(\ln\rho) + \zeta(e_r)B_2(e_r, e_s, Je_j) = \rho e_r^0(B_2^0(e_r^0, e_s^0, e_j^0)) \\ &\quad + e_s(e_j(\ln\rho)) - \rho B_2^0(\mathcal{V}\text{grad}_{g_0}\ln(\rho^2\sigma^2\lambda_0^2), e_s^0, e_j^0) \\ &\quad - e_s(\ln(\rho^2\sigma^2\lambda_0^2))e_j(\ln\rho) + 2e_s(\ln\rho)(\mu_0^\flat(e_j) + e_j(\ln\rho)) \\ &\quad + \frac{\sigma^2}{\rho^2}\zeta_0(e_r)(B_2^0(e_r^0, e_s^0, Je_j) + \delta_{rs}Je_j(\ln\rho)) = \rho e_r^0(B_2^0(e_r^0, e_s^0, e_j^0)) \\ &\quad + e_s(e_j(\ln\rho)) - \rho B_2^0(\mathcal{V}\text{grad}_{g_0}\ln(\rho^2\sigma^2\lambda_0^2), e_s^0, e_j^0) - e_s(\ln(\sigma^2\lambda_0^2))e_j(\ln\rho) \\ &\quad + 2e_s(\ln\rho)\mu_0^\flat(e_j) + \frac{\sigma^2}{\rho^2}B_2^0(\zeta_0^\sharp, e_s, Je_j) + \frac{\sigma^2}{\rho^2}\zeta_0(e_s)Je_j(\ln\rho) \\ &= \text{div}_{g_0}B_2^0(e_s, e_j) + B_2^0(\mathcal{V}\nabla_{e_a^0}e_a^0, e_s, e_j) + B_2^0(e_r^0, e_s, \mathcal{H}\nabla_{e_r^0}e_j) \\ &\quad + e_s(e_j(\ln\rho)) - \rho B_2^0(\mathcal{V}\text{grad}_{g_0}\ln(\rho^2\sigma^2\lambda_0^2), e_s^0, e_j^0) \\ &\quad - e_s(\ln(\sigma^2\lambda_0^2))e_j(\ln\rho) + 2e_s(\ln\rho)\mu_0^\flat(e_j) + \frac{\sigma^2}{\rho^2}B_2^0(\zeta_0^\sharp, e_s, Je_j) \\ &\quad + \frac{\sigma^2}{\rho^2}\zeta_0(e_s)Je_j(\ln\rho) = \text{div}_{g_0}B_2^0(e_s, e_j) + B_2^0(\mathcal{V}\nabla_{e_a^0}e_a^0, e_s, e_j) \\ &\quad + B_2^0(e_r^0, e_s, -\zeta_0(e_r^0)Je_j + e_r^0(\ln\sigma)e_j) + e_s(e_j(\ln\rho)) \\ &\quad - B_2^0(\mathcal{V}\text{grad}_{g_0}\ln(\rho^2\sigma^2\lambda_0^2), e_s, e_j) - e_s(\ln(\sigma^2\lambda_0^2))e_j(\ln\rho) \\ &\quad + 2e_s(\ln\rho)\mu_0^\flat(e_j) + \frac{\sigma^2}{\rho^2}B_2^0(\zeta_0^\sharp, e_s, Je_j) + \frac{\sigma^2}{\rho^2}\zeta_0(e_s)Je_j(\ln\rho) \\ &= \text{div}_{g_0}B_2^0(e_s, e_j) + 2B_2^0(\mathcal{V}\text{grad}_{g_0}\ln\lambda_0, e_s, e_j) - B_2^0(\zeta_0^\sharp, e_s, Je_j) \\ &\quad + B_2^0(\mathcal{V}\text{grad}_{g_0}\ln\sigma, e_s, e_j) + e_s(e_j(\ln\rho)) \\ &\quad - B_2^0(\mathcal{V}\text{grad}_{g_0}\ln(\rho^2\sigma^2\lambda_0^2), e_s, e_j) - e_s(\ln(\sigma^2\lambda_0^2))e_j(\ln\rho) \\ &\quad + 2e_s(\ln\rho)\mu_0^\flat(e_j) + \frac{\sigma^2}{\rho^2}B_2^0(\zeta_0^\sharp, e_s, Je_j) + \frac{\sigma^2}{\rho^2}\zeta_0(e_s)Je_j(\ln\rho) \\ &= \text{div}_{g_0}B_2^0(e_s, e_j) + e_s(e_j(\ln\rho)) - B_2^0(\mathcal{V}\text{grad}_{g_0}\ln(\rho^2\sigma), e_s, e_j) \\ &\quad - e_s(\ln(\sigma^2\lambda_0^2))e_j(\ln\rho) + 2e_s(\ln\rho)\mu_0^\flat(e_j) \\ &\quad + \left(\frac{\sigma^2}{\rho^2} - 1\right)B_2^0(\zeta_0^\sharp, e_s, Je_j) + \frac{\sigma^2}{\rho^2}\zeta_0(e_s)Je_j(\ln\rho). \end{aligned}$$

However

$$\begin{aligned}
 \nabla^0 d \ln \rho(e_s, e_j) &= -(\nabla_{e_s}^0 e_j)(\ln \rho) + e_s(e_j(\ln \rho)) \\
 &= -(\mathcal{H} \nabla_{e_s}^0 e_j)(\ln \rho) - (\mathcal{V} \nabla_{e_s}^0 e_j)(\ln \rho) + e_s(e_j(\ln \rho)) \\
 &= \zeta_0(e_s) J e_j(\ln \rho) - e_s(\ln \sigma) e_j(\ln \rho) + B_{e_s}^{0*} e_j(\ln \rho) + e_s(e_j(\ln \rho)).
 \end{aligned}$$

Finally,

$$\begin{aligned}
 B_{e_s}^{0*} e_j(\ln \rho) &= g_0(\mathcal{V} \text{grad} \ln \rho, B_{e_s}^{0*} e_j) = g_0(B_{e_s}^0 \mathcal{V} \text{grad} \ln \rho, e_j) \\
 &= B_2^0(\mathcal{V} \text{grad} \ln \rho, e_s, e_j)
 \end{aligned}$$

and the expression follows. \square

Lemma 4.14. *Under biconformal deformation, the quantities C and C^* change according to*

$$\begin{aligned}
 C &= \frac{\sigma^2}{\rho^2} \{C_0 + d \ln \rho(B_{\star}^0 \star) + \|\mathcal{H} \text{grad}_{g_0} \ln \rho\|_{g_0}^2 g_0^{\mathcal{V}}\} \\
 C^* &= C_0^* + 4d \ln \rho \odot \mu_0^b + 2(d \ln \rho \circ \mathcal{H})^2.
 \end{aligned}$$

Proof. From Corollary (4.3),

$$\begin{aligned}
 C(e_r, e_s) &= g(B_{e_t} e_r, B_{e_t} e_s) = \frac{1}{\sigma^2} g_0(B_{e_t} e_r, B_{e_t} e_s) \\
 &= \frac{1}{\sigma^2} g_0(\sigma^2(B_{e_t}^0 e_r^0 + \delta_{rt} \mathcal{H} \text{grad}_{g_0} \ln \rho), \sigma^2(B_{e_t}^0 e_s^0 + \delta_{st} \mathcal{H} \text{grad}_{g_0} \ln \rho)) \\
 &= \frac{\sigma^2}{\rho^2} C_0(e_r, e_s) + \frac{2\sigma^2}{\rho^2} d \ln \rho(B_{e_r}^0 e_s) + \frac{\sigma^2}{\rho^2} \|\mathcal{H} \text{grad}_{g_0} \ln \rho\|_{g_0}^2 g_0(e_r, e_s),
 \end{aligned}$$

whereas

$$\begin{aligned}
 C^*(e_i, e_j) &= g(B_{e_r}^* e_i, B_{e_r}^* e_j) = g(e_s, B_{e_r}^* e_i) g(e_s, B_{e_r}^* e_j) = g(B_{e_r} e_s, e_i) g(B_{e_r} e_s, e_j) \\
 &= \frac{1}{\sigma^4} g_0(\sigma^2(B_{e_r}^0 e_s^0 + \delta_{rs} \mathcal{H} \text{grad}_{g_0} \ln \rho), e_i) \\
 &\quad \times g_0(\sigma^2(B_{e_r}^0 e_s^0 + \delta_{rs} \mathcal{H} \text{grad}_{g_0} \ln \rho), e_j) \\
 &= C_0^*(e_i, e_j) + 2d \ln \rho(e_i) \mu_0^b(e_j) + 2d \ln \rho(e_j) \mu_0^b(e_i) + 2d \ln \rho(e_i) d \ln \rho(e_j).
 \end{aligned}$$

\square

Remark 4.15. When $\sigma = \rho$, the deformation is conformal and there is a well-known formula for the change in Ricci [7]:

$$\begin{aligned}
 \text{Ric}(e_a, e_b) &= \text{Ric}^0(e_a, e_b) + 2[\nabla^0 d \ln \sigma(e_a, e_b) + e_a(\ln \sigma) e_b(\ln \sigma)] \\
 &\quad + (\Delta_{g_0} \ln \sigma - 2\|\text{grad}_{g_0} \ln \sigma\|^2) g_0(e_a, e_b).
 \end{aligned}$$

5. ORTHOGONAL PROJECTION FROM \mathbb{R}^4 TO \mathbb{R}^2

Let $\varphi: \mathbb{R}^4 \rightarrow \mathbb{R}^2$ be the canonical projection $\varphi(x^1, x^2, x^3, x^4) = (x^1, x^2)$. Then $\lambda_0 \equiv 1, \mu_0 \equiv 0, B^0 = B_1^0 = B_2^0 \equiv 0, \zeta^0 \equiv 0$. We take the standard basis: $e_a^0 = \partial/\partial x^a$.

From Lemma 3.4,

$$\begin{aligned} \text{Ric}|_{H \times H} &= \{\lambda^2 K^N + \Delta \ln \lambda + 2d \ln \lambda(\mu) - 2\|\zeta\|^2\} g^{\mathcal{H}} - C^* + \mathcal{L}_\mu g|_{H \times H} \\ &= \{\Delta \ln \lambda + 2d \ln \lambda(\mu)\} g^{\mathcal{H}} - C^* + \mathcal{L}_\mu g|_{H \times H}, \end{aligned}$$

where $\lambda = \sigma$ and $\mu = \sigma^2 \mathcal{H} \text{grad}_{g_0} \ln \rho$.

From Corollary 4.9,

$$\begin{aligned} \Delta_g \ln \lambda &= \sigma^2 \Delta_{g_0} \ln \sigma + (\rho^2 - \sigma^2) \Delta_{g_0}^{\mathcal{V}}(\ln \sigma) - 2\sigma^2 d \ln \sigma(\mathcal{H} \text{grad}_{g_0} \ln \rho) \\ &\quad - 2\rho^2 d \ln \sigma(\mathcal{V} \text{grad}_{g_0} \ln \sigma) \end{aligned}$$

and $d \ln \lambda(\mu) = \sigma^2 d \ln \sigma(\mathcal{H} \text{grad}_{g_0} \ln \rho)$, so that

$$\begin{aligned} \Delta_g \ln \lambda + 2d \ln \lambda(\mu) &= \sigma^2 \left(\frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} \right) + \rho^2 \left(\frac{\partial^2 \ln \sigma}{\partial x_3^2} + \frac{\partial^2 \ln \sigma}{\partial x_4^2} \right) \\ &\quad - 2\rho^2 \left(\left(\frac{\partial \ln \sigma}{\partial x_3} \right)^2 + \left(\frac{\partial \ln \sigma}{\partial x_4} \right)^2 \right). \end{aligned}$$

From Lemma 4.14,

$$C^*(e_i, e_j) = 2\sigma^2 e_i^0(\ln \rho) e_j^0(\ln \rho) = 2\sigma^2 \frac{\partial \ln \rho}{\partial x^i} \frac{\partial \ln \rho}{\partial x^j}.$$

Next, from Lemma 4.1(v),

$$\begin{aligned} \mathcal{L}_\mu g(e_i, e_j) &= g(\nabla_{e_i} \mu, e_j) + g(e_i, \nabla_{e_j} \mu) \\ &= e_i(g(\mu, e_j)) + e_j(g(\mu, e_i)) - g(\mu, \nabla_{e_i} e_j + \nabla_{e_j} e_i) \\ &= e_i(e_j(\ln \rho)) + e_j(e_i(\ln \rho)) - e_k(\ln \rho) g(e_k, \nabla_{e_i} e_j + \nabla_{e_j} e_i) \\ &= e_i(e_j(\ln \rho)) + e_j(e_i(\ln \rho)) + e_i(\ln \rho) e_j(\ln \sigma) \\ &\quad + e_i(\ln \sigma) e_j(\ln \rho) - 2\delta_{ij} \mathcal{H} \text{grad}_g \ln \rho \\ &= 2\sigma^2 \left\{ \frac{\partial^2 \ln \rho}{\partial x^i \partial x^j} + \frac{\partial \ln \sigma}{\partial x^i} \frac{\partial \ln \rho}{\partial x^j} + \frac{\partial \ln \rho}{\partial x^i} \frac{\partial \ln \sigma}{\partial x^j} \right. \\ &\quad \left. - \delta_{ij} \left(\frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \rho}{\partial x_1} + \frac{\partial \ln \sigma}{\partial x_2} \frac{\partial \ln \rho}{\partial x_2} \right) \right\}. \end{aligned}$$

Collecting terms, we obtain

$$\begin{aligned}
 \text{Ric}(e_1, e_1) &= \sigma^2 \left\{ \left(\frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} \right) + \frac{\rho^2}{\sigma^2} \left(\frac{\partial^2 \ln \sigma}{\partial x_3^2} + \frac{\partial^2 \ln \sigma}{\partial x_4^2} \right) \right. \\
 &\quad \left. - 2 \frac{\rho^2}{\sigma^2} \left(\left(\frac{\partial \ln \sigma}{\partial x_3} \right)^2 + \left(\frac{\partial \ln \sigma}{\partial x_4} \right)^2 \right) \right. \\
 &\quad \left. - 2 \left(\frac{\partial \ln \rho}{\partial x_1} \right)^2 + 2 \frac{\partial^2 \ln \rho}{\partial x_1^2} + 2 \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \rho}{\partial x_1} - 2 \frac{\partial \ln \sigma}{\partial x_2} \frac{\partial \ln \rho}{\partial x_2} \right\} \\
 \text{Ric}(e_2, e_2) &= \sigma^2 \left\{ \left(\frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} \right) + \frac{\rho^2}{\sigma^2} \left(\frac{\partial^2 \ln \sigma}{\partial x_3^2} + \frac{\partial^2 \ln \sigma}{\partial x_4^2} \right) \right. \\
 &\quad \left. - 2 \frac{\rho^2}{\sigma^2} \left(\left(\frac{\partial \ln \sigma}{\partial x_3} \right)^2 + \left(\frac{\partial \ln \sigma}{\partial x_4} \right)^2 \right) \right. \\
 &\quad \left. - 2 \left(\frac{\partial \ln \rho}{\partial x_2} \right)^2 + 2 \frac{\partial^2 \ln \rho}{\partial x_2^2} + 2 \frac{\partial \ln \sigma}{\partial x_2} \frac{\partial \ln \rho}{\partial x_2} - 2 \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \rho}{\partial x_1} \right\} \\
 \text{Ric}(e_1, e_2) &= 2\sigma^2 \left\{ \frac{\partial^2 \ln \rho}{\partial x_1 \partial x_2} - \frac{\partial \ln \rho}{\partial x_1} \frac{\partial \ln \rho}{\partial x_2} + \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \rho}{\partial x_2} + \frac{\partial \ln \rho}{\partial x_1} \frac{\partial \ln \sigma}{\partial x_2} \right\}.
 \end{aligned}$$

From Lemma 3.5, the mixed Ricci tensor acting on (e_j, e_s) is given by

$$\begin{aligned}
 \text{Ric}(e_j, e_s) &= \nabla d \ln \lambda(e_j, e_s) - (d \ln \lambda)^2(e_j, e_s) - 2(d \ln \lambda \odot \zeta)(Je_j, e_s) \\
 &\quad - (\nabla_{Je_j} \zeta)(e_s) - 2\zeta(\nabla_{e_s} Je_j) \\
 &\quad - \text{div } B_2(e_s, e_j) - 2d \ln \lambda(B_{e_s}^* e_j) + 2(\nabla_{e_s} \mu^b)(e_j) \\
 &= \nabla d \ln \sigma(e_j, e_s) - (d \ln \sigma)^2(e_j, e_s) - \text{div } B_2(e_s, e_j) \\
 &\quad - 2d \ln \lambda(B_{e_s}^* e_j) + 2(\nabla_{e_s} \mu^b)(e_j).
 \end{aligned}$$

From Corollary 4.2,

$$\nabla d \ln \sigma(e_s, e_j) = e_s(e_j(\ln \sigma)) - d \ln \sigma(\nabla_{e_s} e_j) = e_s(e_j(\ln \sigma)) + e_j(\ln \rho)e_s(\ln \sigma).$$

Since the fibres before deformation are totally geodesic, $B^0 \equiv 0$, so from Lemma 4.13,

$$\begin{aligned}
 (\text{div } B_2)(e_s, e_j) &= \nabla^0 d \ln \rho(e_s, e_j) - e_s(\ln \sigma)e_j(\ln \rho) = e_s(e_j(\ln \rho)) \\
 &\quad - d \ln \rho(\nabla_{e_s}^0 e_j) - e_s(\ln \sigma)e_j(\ln \rho) \\
 &= e_s(e_j(\ln \rho)) - d \ln \rho(\sigma \rho \nabla_{e_s}^0 e_j^0 + \sigma \rho e_s^0(\ln \sigma)e_j^0) - e_s(\ln \sigma)e_j(\ln \rho) \\
 &= e_s(e_j(\ln \rho)) - 2e_s(\ln \sigma)e_j(\ln \rho).
 \end{aligned}$$

From Corollary (4.2),

$$d \ln \lambda(B_{e_s}^* e_j) = -d \ln \sigma(\mathcal{V} \nabla_{e_s} e_j) = -d \ln \sigma(e_r)g(e_r, \nabla_{e_s} e_j) = e_s(\ln \sigma)e_j(\ln \rho).$$

Finally, $\mu = \sigma^2 \mathcal{H} \text{grad}_{g_0} \ln \rho = e_i(\ln \rho)e_i$, so that from Corollary 4.2,

$$\begin{aligned}
 (\nabla_{e_s} \mu^b)(e_j) &= e_s(g(\mu, e_j)) - g(\mu, \nabla_{e_s} e_j) \\
 &= e_s(e_i(\ln \rho)\delta_{ij}) - e_i(\ln \rho)g(e_i, \nabla_{e_s} e_j) = e_s(e_j(\ln \rho)).
 \end{aligned}$$

We conclude that

$$\text{Ric}(e_j, e_s) = e_s(e_j(\ln \sigma)) + e_s(e_j(\ln \rho)) + e_j(\ln \rho)e_s(\ln \sigma) - e_j(\ln \sigma)e_s(\ln \sigma),$$

explicitly

$$\text{Ric}(e_j, e_s) = \sigma \rho \left\{ \frac{\partial^2 \ln(\sigma \rho)}{\partial x^j \partial x^s} + 2 \frac{\partial \ln \sigma}{\partial x^s} \frac{\partial \ln \rho}{\partial x^j} \right\}.$$

The vertical components of the Ricci tensor are given by

$$\begin{aligned} \text{Ric}|_{V \times V} &= K^V g^V + 2 \nabla \text{d} \ln \lambda|_{V \times V} + 2 \text{d} \ln \lambda (B_\star \star) - 2 (\text{d} \ln \lambda)^2|_{V \times V} \\ &\quad + 2 \zeta^2 + \text{div } B_1|_{V \times V} \\ &= K^V g^V + 2 \nabla \text{d} \ln \lambda|_{V \times V} + 2 \text{d} \ln \lambda (B_\star \star) - 2 (\text{d} \ln \lambda)^2|_{V \times V} + \text{div } B_1|_{V \times V}. \end{aligned}$$

After biconformal deformation, the sectional curvature of the fibres is given by

$$K^V = \rho^2 \Delta_{g_0}^V \ln \rho.$$

For the second fundamental form:

$$\nabla \text{d} \ln \lambda(e_r, e_s) = e_r(e_s(\ln \lambda)) - \text{d} \ln \lambda(\nabla_{e_r} e_s).$$

From Corollary 4.2,

$$\nabla_{e_r} e_s = \delta_{rs} \{ \sigma^2 \mathcal{H} \text{grad}_{g_0} \ln \rho + \rho^2 \mathcal{V} \text{grad}_{g_0} \ln \rho \} - e_s(\ln \rho) e_r$$

and

$$\begin{aligned} \nabla \text{d} \ln \lambda(e_r, e_s) &= e_r(e_s(\ln \lambda)) - \text{d} \ln \lambda(\nabla_{e_r} e_s) \\ &= \rho^2 e_r^0(e_s^0(\ln \sigma)) + \rho^2 e_r^0(\ln \rho) e_s^0(\ln \sigma) + \rho^2 e_s^0(\ln \rho) e_r^0(\ln \sigma) \\ &\quad - \delta_{rs} \text{d} \ln \sigma (\sigma^2 \mathcal{H} \text{grad}_{g_0} \ln \rho + \rho^2 \mathcal{V} \text{grad}_{g_0} \ln \rho) \end{aligned}$$

From Corollary (4.3),

$$B_{e_r} e_s = \sigma^2 \delta_{rs} \mathcal{H} \text{grad}_{g_0} \ln \rho.$$

From Lemma 4.11 we have

$$(\text{div } B_1)(e_r, e_s) = \delta_{rs} \sigma^2 \{ \text{Tr}_{g_0}^{\mathcal{H}} \nabla \text{d} \ln \rho - \text{d} \ln \rho (\mathcal{H} \text{grad}_{g_0} \ln \rho^2) \}.$$

Thus

$$\begin{aligned} \text{Ric}(e_r, e_s) &= \rho^2 \delta_{rs} \Delta_{g_0}^V \ln \rho - 2 \rho^2 e_r^0(\ln \sigma) e_s^0(\ln \sigma) \\ &\quad + 2 \{ \rho^2 e_r^0(e_s^0(\ln \sigma)) + \rho^2 e_r^0(\ln \rho) e_s^0(\ln \sigma) + \rho^2 e_s^0(\ln \rho) e_r^0(\ln \sigma) \} \\ &\quad + \delta_{rs} \{ \sigma^2 \text{Tr}_{g_0}^{\mathcal{H}} \nabla \text{d} \ln \rho - 2 \sigma^2 \text{d} \ln \rho (\mathcal{H} \text{grad}_{g_0} \ln \rho) \\ &\quad - 2 \rho^2 \text{d} \ln \sigma (\mathcal{V} \text{grad}_{g_0} \ln \rho) \}. \end{aligned}$$

Explicitly,

$$\begin{aligned} \text{Ric}(e_r, e_s) = & \rho^2 \left\{ 2 \frac{\partial^2 \ln \sigma}{\partial x^r \partial x^s} + 2 \frac{\partial \ln \rho}{\partial x^r} \frac{\partial \ln \sigma}{\partial x^s} + 2 \frac{\partial \ln \sigma}{\partial x^r} \frac{\partial \ln \rho}{\partial x^s} - 2 \frac{\partial \ln \sigma}{\partial x^r} \frac{\partial \ln \sigma}{\partial x^s} \right. \\ & + \delta_{rs} \left(\frac{\sigma^2}{\rho^2} \left(\frac{\partial^2 \ln \rho}{\partial x_1^2} + \frac{\partial^2 \ln \rho}{\partial x_2^2} \right) + \frac{\partial^2 \ln \rho}{\partial x_3^2} + \frac{\partial^2 \ln \rho}{\partial x_4^2} \right. \\ & \left. \left. - 2 \frac{\sigma^2}{\rho^2} \left(\left(\frac{\partial \ln \rho}{\partial x_1} \right)^2 + \left(\frac{\partial \ln \rho}{\partial x_2} \right)^2 \right) - 2 \left(\frac{\partial \ln \sigma}{\partial x_3} \frac{\partial \ln \rho}{\partial x_3} + \frac{\partial \ln \sigma}{\partial x_4} \frac{\partial \ln \rho}{\partial x_4} \right) \right) \right\}. \end{aligned}$$

The equations for an Einstein metric: $\text{Ric} = Ag$ for some constant A , become the following system of ten equations:

(2)

$$\begin{aligned} \text{(i)} \quad A = & \sigma^2 \left\{ \frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} + \frac{\rho^2}{\sigma^2} \left(\frac{\partial^2 \ln \sigma}{\partial x_3^2} + \frac{\partial^2 \ln \sigma}{\partial x_4^2} \right) \right. \\ & - 2 \frac{\rho^2}{\sigma^2} \left(\left(\frac{\partial \ln \sigma}{\partial x_3} \right)^2 + \left(\frac{\partial \ln \sigma}{\partial x_4} \right)^2 \right) + 2 \frac{\partial^2 \ln \rho}{\partial x_j^2} \\ & \left. - 2 \left(\frac{\partial \ln \rho}{\partial x_j} \right)^2 + 2 \frac{\partial \ln \sigma}{\partial x_j} \frac{\partial \ln \rho}{\partial x_j} - 2 \frac{\partial \ln \sigma}{\partial x_{j'}} \frac{\partial \ln \rho}{\partial x_{j'}} \right\} \quad (j = 1, 2) \\ \text{(ii)} \quad 0 = & \frac{\partial^2 \ln \rho}{\partial x_1 \partial x_2} + \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \rho}{\partial x_2} + \frac{\partial \ln \rho}{\partial x_1} \frac{\partial \ln \sigma}{\partial x_2} - \frac{\partial \ln \rho}{\partial x_1} \frac{\partial \ln \rho}{\partial x_2} \\ \text{(iii)} \quad 0 = & \frac{\partial^2 \ln(\sigma \rho)}{\partial x^j \partial x^s} + 2 \frac{\partial \ln \sigma}{\partial x^s} \frac{\partial \ln \rho}{\partial x^j} \quad (j = 1, 2, s = 3, 4) \\ \text{(iv)} \quad A = & \rho^2 \left\{ 2 \frac{\partial^2 \ln \sigma}{\partial x_s^2} - 2 \left(\frac{\partial \ln \sigma}{\partial x_s} \right)^2 + 2 \frac{\partial \ln \sigma}{\partial x_s} \frac{\partial \ln \rho}{\partial x_s} - 2 \frac{\partial \ln \sigma}{\partial x_{s'}} \frac{\partial \ln \rho}{\partial x_{s'}} \right. \\ & + \frac{\sigma^2}{\rho^2} \left(\frac{\partial^2 \ln \rho}{\partial x_1^2} + \frac{\partial^2 \ln \rho}{\partial x_2^2} \right) + \frac{\partial^2 \ln \rho}{\partial x_3^2} + \frac{\partial^2 \ln \rho}{\partial x_4^2} \\ & \left. - 2 \frac{\sigma^2}{\rho^2} \left(\left(\frac{\partial \ln \rho}{\partial x_1} \right)^2 + \left(\frac{\partial \ln \rho}{\partial x_2} \right)^2 \right) \right\} \quad (s = 3, 4) \\ \text{(v)} \quad 0 = & \frac{\partial^2 \ln \sigma}{\partial x_3 \partial x_4} + \frac{\partial \ln \rho}{\partial x_3} \frac{\partial \ln \sigma}{\partial x_4} + \frac{\partial \ln \sigma}{\partial x_3} \frac{\partial \ln \rho}{\partial x_4} - \frac{\partial \ln \sigma}{\partial x_3} \frac{\partial \ln \sigma}{\partial x_4}. \end{aligned}$$

Note the symmetry between the equations: after the interchange $(j, k, \sigma, \rho) \leftrightarrow (r, s, \rho, \sigma)$, equations (i) and (iv) are interchanged.

5.1. Warped product solutions. Let us investigate some special solutions. If $\sigma = \sigma(x_1, x_2)$ and $\rho = \rho(x_3, x_4)$, then the system reduces to

$$\frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} = A/\sigma^2 \quad \text{and} \quad \frac{\partial^2 \ln \rho}{\partial x_3^2} + \frac{\partial^2 \ln \rho}{\partial x_4^2} = A/\rho^2.$$

Note that $A = \sigma^2 \left(\frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} \right)$ is the Gaussian curvature of the surface $(\mathbb{R}^2, (dx_1^2 + dx_2^2)/\sigma^2)$; similarly for the second equation. For example, setting

$$\sigma = \frac{1 + x_1^2 + x_2^2}{2} \quad \text{and} \quad \rho = \frac{1 + x_3^2 + x_4^2}{2}$$

yields the product of spheres $S^2 \times S^2$ with constant $A = 1$, whereas setting

$$\sigma = \frac{1 - x_1^2 - x_2^2}{2} \quad \text{and} \quad \rho = \frac{1 - x_3^2 - x_4^2}{2}$$

yields the product of hyperbolic spaces $H^2 \times H^2$ with constant $A = -1$.

More generally a warped product of the surfaces $(\mathbb{R}^2, (dx_1^2 + dx_2^2)/\sigma(x_1, x_2)^2)$ and $(\mathbb{R}^2, (dx_3^2 + dx_4^2)/\beta(x_3, x_4)^2)$ corresponds to \mathbb{R}^4 endowed with a metric of the form:

$$(3) \quad g = \frac{dx_1^2 + dx_2^2}{\sigma(x_1, x_2)^2} + \frac{dx_3^2 + dx_4^2}{\alpha(x_1, x_2)^2 \beta(x_3, x_4)^2}.$$

Setting $\sigma = \sigma(x_1, x_2)$ and $\rho = \alpha(x_1, x_2)\beta(x_3, x_4)$, the Einstein equations become the system:

$$(4) \quad \begin{aligned} (i) \quad A &= \sigma^2 \left\{ \frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} - 2 \left(\frac{\partial \ln \alpha}{\partial x_1} \right)^2 + 2 \frac{\partial^2 \ln \alpha}{\partial x_1^2} \right. \\ &\quad \left. + 2 \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \alpha}{\partial x_1} - 2 \frac{\partial \ln \sigma}{\partial x_2} \frac{\partial \ln \alpha}{\partial x_2} \right\} \\ (ii) \quad A &= \sigma^2 \left\{ \frac{\partial^2 \ln \sigma}{\partial x_1^2} + \frac{\partial^2 \ln \sigma}{\partial x_2^2} - 2 \left(\frac{\partial \ln \alpha}{\partial x_2} \right)^2 + 2 \frac{\partial^2 \ln \alpha}{\partial x_2^2} \right. \\ &\quad \left. + 2 \frac{\partial \ln \sigma}{\partial x_2} \frac{\partial \ln \alpha}{\partial x_2} - 2 \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \alpha}{\partial x_1} \right\} \\ (iii) \quad 0 &= \frac{\partial^2 \ln \alpha}{\partial x_1 \partial x_2} + \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \alpha}{\partial x_2} + \frac{\partial \ln \alpha}{\partial x_1} \frac{\partial \ln \sigma}{\partial x_2} - \frac{\partial \ln \alpha}{\partial x_1} \frac{\partial \ln \alpha}{\partial x_2} \\ (iv) \quad A &= \sigma^2 \left(\frac{\partial^2 \ln \alpha}{\partial x_1^2} + \frac{\partial^2 \ln \alpha}{\partial x_2^2} \right) + \alpha^2 \beta^2 \left(\frac{\partial^2 \ln \beta}{\partial x_3^2} + \frac{\partial^2 \ln \beta}{\partial x_4^2} \right) \\ &\quad - 2\sigma^2 \left(\left(\frac{\partial \ln \alpha}{\partial x_1} \right)^2 + \left(\frac{\partial \ln \alpha}{\partial x_2} \right)^2 \right). \end{aligned}$$

The sum of (i) and (ii) gives the equation:

$$A = \sigma^2 (\Delta_{g_0} \ln \sigma + \Delta_{g_0} \ln \alpha - \|\text{grad}_{g_0} \ln \alpha\|_0^2).$$

On the other hand the difference gives:

$$0 = \frac{\partial^2 \ln \alpha}{\partial x_1^2} - \frac{\partial^2 \ln \alpha}{\partial x_2^2} - \left(\frac{\partial \ln \alpha}{\partial x_1} \right)^2 + \left(\frac{\partial \ln \alpha}{\partial x_2} \right)^2 + 2 \frac{\partial \ln \sigma}{\partial x_1} \frac{\partial \ln \alpha}{\partial x_1} - 2 \frac{\partial \ln \sigma}{\partial x_2} \frac{\partial \ln \alpha}{\partial x_2}.$$

Since $\alpha = \alpha(x_1, x_2)$ and $\beta = \beta(x_3, x_4)$ are independent, on dividing equation (iv) by α^2 , we deduce that

$$(5) \quad \beta^2 \left(\frac{\partial^2 \ln \beta}{\partial x_3^2} + \frac{\partial^2 \ln \beta}{\partial x_4^2} \right) = C$$

for a constant C and in particular the metric $(dx_3^2 + dx_4^2)/\beta^2$ is necessarily of constant Gaussian curvature C , and

$$A - C\alpha^2 = \sigma^2 \Delta_{g_0} \ln \alpha - 2\sigma^2 \|\text{grad}_{g_0} \ln \alpha\|_0^2.$$

Set $x_1 = t$ and suppose that $\alpha = \alpha(t)$ and $\sigma = \sigma(t)$ depend only on t . Then (4)(iii) is satisfied and α and σ are determined by the system:

$$(6) \quad \begin{cases} \text{(i)} & A = \sigma^2 ((\ln \sigma)'' + (\ln \alpha)'' - (\ln \alpha)'^2) \\ \text{(ii)} & 0 = (\ln \alpha)'' - (\ln \alpha)'^2 + 2(\ln \sigma)'(\ln \alpha)' \\ \text{(iii)} & A - C\alpha^2 = \sigma^2 ((\ln \alpha)'' - 2(\ln \alpha)'^2) \end{cases}.$$

From (6)(ii), provided $(\ln \alpha)' \neq 0$,

$$\begin{aligned} 2(\ln \sigma)' &= \frac{-(\ln \alpha)'' + (\ln \alpha)'^2}{(\ln \alpha)'} = (-\ln |(\ln \alpha)'| + \ln \alpha)' \\ \implies 2 \ln \sigma &= -\ln |(\ln \alpha)'| + \ln \alpha + a \implies \sigma^2 = B\alpha^2/\alpha', \end{aligned}$$

for constants a and B , with B non-zero. In particular, taking the difference between (6)(i) and (iii), we deduce that

$$\tilde{C}\alpha' = (\ln \sigma)'' + (\ln \alpha)'^2,$$

where $\tilde{C} = C/B$. But from (6)(ii),

$$\begin{aligned} 2(\ln \sigma)'' &= \frac{-(\ln \alpha)''' + (\ln \alpha)'(\ln \alpha)''}{(\ln \alpha)'} + \frac{(\ln \alpha)''^2}{(\ln \alpha)'^2} \\ \implies 2\tilde{C}\alpha' &= \frac{-(\ln \alpha)''' + (\ln \alpha)'(\ln \alpha)''}{(\ln \alpha)'} + \frac{(\ln \alpha)''^2}{(\ln \alpha)'^2} + 2(\ln \alpha)'^2. \end{aligned}$$

This simplifies to the third order ODE:

$$\alpha\alpha''' = 2\alpha'\alpha'' + \frac{\alpha(\alpha'')^2}{\alpha'} - 2\tilde{C}\alpha\alpha'^2.$$

Note the specific solution $\alpha(t) = t$ corresponding to hyperbolic space. More generally, if we set $\gamma(t) = \alpha'(t)$, $\delta(t) = \alpha''(t) = \gamma'(t)$, then we have the first order system:

$$(7) \quad \begin{pmatrix} \alpha \\ \gamma \\ \delta \end{pmatrix}' = \begin{pmatrix} \gamma \\ \delta \\ \frac{2\gamma\delta}{\alpha} + \frac{\delta^2}{\gamma} - 2\tilde{C}\gamma^2 \end{pmatrix}.$$

Cauchy's existence theorem (see, for example, [5] (10.4.5)) yields local solutions:

Let $\Gamma_0 = \begin{pmatrix} \alpha_0 \\ \gamma_0 \\ \delta_0 \end{pmatrix} \in \mathbb{R}^3$ be a point with $\alpha_0 > 0$ and $\gamma_0 \neq 0$ and let $t_0 \in \mathbb{R}$. Then

there is a solution $\Gamma(t) = \begin{pmatrix} \alpha(t) \\ \gamma(t) \\ \delta(t) \end{pmatrix}$ to (7) defined on an open interval $I \subset \mathbb{R}$ ($t_0 \in I$), with $\Gamma(t_0) = \Gamma_0$.

Given such a solution to (7) on an open interval I with $\alpha(t)$ positive and $\alpha'(t)$ non-zero for all $t \in I$, then defining σ by $\sigma^2 = B\alpha^2/\alpha'$, where B is a non-zero constant of sign consistent with α' and where we require $C = B\tilde{C}$ to be the constant Gaussian curvature of the metric $(dx_3^2 + dx_4^2)/\beta(x_3, x_4)^2$, equations (6) are satisfied and the metric (3) is Einstein. The constant A is given by (6)(iii):

$$A = C\alpha^2 + \frac{B\alpha^2}{\alpha'} \left(\frac{\alpha''}{\alpha} - \frac{3\alpha'^2}{\alpha^2} \right),$$

which one easily checks is an integral of (7).

5.2. Solutions depending on a single parameter. Replace x_1 with the parameter t and suppose that both σ and ρ depend only on t . Then (2)(iii), (iv) and (vii) are satisfied, while (i) becomes

$$(8) \quad A = \sigma^2 \{ (\ln \sigma)'' + 2(\ln \sigma)'(\ln \rho)' - 2(\ln \rho)'^2 + 2(\ln \rho)'' \};$$

(ii) becomes

$$(9) \quad A = \sigma^2 \{ (\ln \sigma)'' - 2(\ln \sigma)'(\ln \rho)' \};$$

(v) and (vi) become

$$(10) \quad A = \sigma^2 \{ (\ln \rho)'' - 2(\ln \rho)'^2 \}.$$

The first two of these are equivalent to the pair of equations:

$$(11) \quad \begin{cases} (a) & A = \sigma^2 (\ln \sigma)'' - 2\sigma^2 (\ln \sigma)'(\ln \rho)' \\ (b) & 0 = -(\ln \rho)'^2 + (\ln \rho)'' + 2(\ln \sigma)'(\ln \rho)' \end{cases}$$

while the third becomes

$$(12) \quad \begin{aligned} A = \sigma^2 (\ln \rho)'' - 2\sigma^2 (\ln \rho)'^2 & \stackrel{(b)}{\implies} \frac{A}{\sigma^2} = -(\ln \rho)'^2 - 2(\ln \sigma)'(\ln \rho)' \\ & \stackrel{(a)}{\implies} -(\ln \rho)'^2 = (\ln \sigma)'' . \end{aligned}$$

We can combine (11)(a) and the first identity of (12) to deduce

$$\begin{aligned} (\ln \rho)'' - (\ln \sigma)'' &= 2(\ln \rho)'((\ln \rho)' - (\ln \sigma)') \Rightarrow \left(\ln \left(\frac{\rho}{\sigma} \right) \right)'' = 2(\ln \rho)' \left(\ln \left(\frac{\rho}{\sigma} \right) \right)' \\ &\Rightarrow (\ln |(\ln u)'|)' = 2(\ln \rho)' \Rightarrow (\ln u)' = c\rho^2 \end{aligned}$$

for a constant c , where we have written $u = \rho/\sigma$. This determines σ as a function of ρ :

$$(13) \quad \frac{\rho}{\sigma} = (1/a)e^{\int c\rho^2 dt} \Rightarrow \sigma = a\rho e^{-\int c\rho^2 dt}$$

for constants a and c . It also yields the identity:

$$(\ln \rho)' - (\ln \sigma)' = c\rho^2 \implies (\ln \sigma)'' = (\ln \rho)'' - 2c\rho\rho'.$$

When we combine this with the last identity of (12), we obtain

$$(14) \quad \begin{aligned} (\ln \rho)'' + (\ln \rho)'^2 = 2c\rho\rho' &\implies \rho'' = 2c\rho^2\rho' \\ &\implies \rho' = \frac{2c}{3}\rho^3 + e \end{aligned}$$

for another constant e . Then from (13):

$$(15) \quad \sigma = a\rho e^{-\int c\rho^2 dt} = a\rho e^{-\int \frac{\rho''}{2\rho'} dt} = a\rho e^{-\frac{1}{2} \ln |\rho'| + B} = b\rho |\rho'|^{-1/2},$$

for constants B and b where $A = \begin{cases} -3b^2e & \text{if } \rho' > 0 \\ +3b^2e & \text{if } \rho' < 0 \end{cases}$. Conversely, given a solution ρ to (14) with σ given by (13), equations (8), (9) and (10) are satisfied with $A = \begin{cases} -3b^2e & \text{if } \rho' > 0 \\ +3b^2e & \text{if } \rho' < 0 \end{cases}$. Specifically,

$$(\ln \sigma)' = (\ln \rho)' - \frac{1}{2} \frac{\rho''}{\rho'} = (\ln \rho)' - c\rho^2 \implies (\ln \sigma)'' = (\ln \rho)'' - 2c\rho\rho'$$

and we now substitute.

Explicit solutions can be obtained by solving (14). In the case when $e = 0$, then up to an affine linear change in the t coordinate, the solution is given by $\rho(t) = t^{-1/2}$ with

$$\sigma(t) = at^{-1/2}e^{\frac{3}{4}\int t^{-1} dt} = at^{1/4}.$$

This corresponds to an incomplete Ricci flat ($A = 0$) metric defined on the half-space $t > 0$.

In the case when $e \neq 0$, relabel the constants such that

$$(16) \quad \rho' = \alpha(\rho^3 - \beta^3) = \alpha(\rho - \beta)(\rho^2 + \beta\rho + \beta^2) \quad (c = 3\alpha/2 \text{ and } e = -\alpha\beta^3).$$

Then

$$\frac{d\rho}{\alpha(\rho - \beta)(\rho^2 + \beta\rho + \beta^2)} = dt$$

which can be integrated explicitly.


Lemma 5.1. (i) For $\alpha < 0$ and $\beta > 0$, there is a solution $\rho(t)$ to (16) that exists for all $t \geq 0$, satisfying $\rho(0) = 0$, $\rho'(t) > 0$ for all $t \geq 0$ and $0 < \rho(t) < \beta$ for all $t > 0$. As $t \rightarrow \infty$, $\rho(t) \rightarrow \beta$ and $\rho'(t) \rightarrow 0$.

(ii) For $\alpha > 0$ and $\beta < 0$, there is a $t_0 > 0$ and a solution $\rho(t)$ to (16) that exists for all $t \in [0, t_0)$ satisfying $\rho(0) = 0$, $\rho'(t) > 0$ for all $t \in [0, t_0)$ and that tends to infinity as $t \rightarrow t_0^-$.

Proof. (i) A solution $\rho(t)$ to (16) in a neighbourhood of $t = 0$ satisfying $\rho(0) = 0$ is guaranteed by the general existence theory of ODEs (see for example [5] (10.4.5)). Without loss of generality we can suppose that $\alpha = -1$ and $\beta = 1$ so the equation has the form

$$(17) \quad \rho' = -\rho^3 + 1.$$

Clearly $\rho'(t) > 0$ provided $\rho(t) < 1$. Suppose that $\rho(t)$ achieves the value 1 and let $t_0 > 0$ be the first time for which this occurs. Then from (17), $\rho'(t_0) = 0$. On differentiating (17), we see that $\rho''(t_0) = -3\rho^2(t_0)\rho'(t_0) = 0$, and so on; by

FIG. 1: Solution to (16) with $\alpha = -1$, $\beta = 1$ and $\rho(0) = 0$

recursion all derivatives $\rho^{(n)}(t_0) = 0$. But by analyticity of the solution (see [5] (10.5.3)), this means that $\rho(t) \equiv 1$ for all t , contradicting the initial condition $\rho(0) = 0$. Thus $\rho(t) < 1$ for all $t \geq 0$.

Clearly any interval of existence $[0, t_1)$ can be extended to $t \geq t_1$, so the solution exists for all time $t \geq 0$ with $\rho(t) \rightarrow 1$ and $\rho'(t) \rightarrow 0$ as $t \rightarrow \infty$.

(ii) Without loss of generality, suppose that $\alpha = 1$ and $\beta = -1$, so that (16) takes the form

$$(18) \quad \rho' = \rho^3 + 1.$$

This time we can appeal to the explicit equation determining ρ obtained on integrating (18) with $\rho(0) = 0$:

$$\frac{1}{3} \ln \frac{\rho + 1}{|\rho^2 - \rho + 1|^{1/2}} + \frac{\sqrt{3}}{3} \arctan \left(\frac{2}{\sqrt{3}} \left(\rho - \frac{1}{2} \right) \right) + \frac{\pi\sqrt{3}}{18} = t.$$

Then as $\rho \rightarrow \infty$, the left-hand side approaches $\frac{2\sqrt{3}\pi}{9}$ which yields the upper bound $t_0 = \frac{2\sqrt{3}\pi}{9}$. \square

In the following theorem, we consider *ends* as components of the complement of the set $\varepsilon \leq t \leq 1/\varepsilon$ for ε small.




FIG. 2: Solution to (16) with $\alpha = 1$, $\beta = -1$ and $\rho(0) = 0$

Theorem 5.2. *Solutions to equation (16) yield two families of 4-dimensional Einstein metrics. Each member of the first family is a complete metric defined on the upper half space $t > 0$, having negative Ricci curvature and two ends: one asymptotic to hyperbolic 4-space and the other to \mathbb{R}^2 . Each member of the second family is incomplete, defined on the space $0 < t < t_0$ for a fixed constant t_0 , and has negative Ricci curvature.*

Proof. Consider the solutions to (16) given by Lemma 5.1(i) and as above, set $e = -\alpha\beta^3 > 0$. At $t = 0$, $\rho(0) = 0$, $\rho'(0) = e$, $\rho''(0) = \rho'''(0) = 0$. Thus the Taylor expansion about $t = 0$ has the form $\rho(t) = et + \mathcal{O}(t^4)$. For σ we have $\sigma(0) = 0$ and

$$\sigma = \frac{b\rho}{\sqrt{\rho'}} \Rightarrow \sigma' = \frac{b(\rho')^{3/2} - \frac{1}{2}b\rho(\rho')^{-1/2}\rho''}{\rho'} \Rightarrow \sigma'(0) = b\sqrt{e}.$$

Furthermore, $\sigma''(0) = \sigma'''(0) = 0$, so that about $t = 0$, we have $\sigma(t) = b\sqrt{e}t + \mathcal{O}(t^4)$. In particular, being of type $g_H := (dt^2 + dx_2^2 + dx_3^2 + dx_4^2)/t^2$, for $t > 0$, the metric is complete in a neighbourhood of the boundary $t = 0$.

The Einstein constant can be deduced from (10), (16) and the expression (15) for σ , specifically $A = 3b^2\alpha\beta^3 < 0$.

In order to study the ends of the resulting Einstein manifold, we consider the exterior to the set $\varepsilon \leq t \leq 1/\varepsilon$ for ε small. As $t \rightarrow \infty$, then $\rho(t) \rightarrow \beta$, $\rho'(t) \rightarrow 0$ and $\sigma(t) \rightarrow \infty$. Thus the metric approaches an end of the form \mathbb{R}^2 with metric $(dx_3^2 + dx_4^2)/\beta^2$. Finally, the Taylor expansions of $\rho(t)$ and $\sigma(t)$ about $t = 0$ show that $g_H - g \rightarrow 0$ as $t \rightarrow 0^+$ (incorporating the constants into

the coordinates), for example $\sigma(t)^2 = t^2 + \mathcal{O}(t^5)$ and $\frac{dt^2 + dx_1^2}{t^2 + \mathcal{O}(t^5)} - \frac{dt^2 + dx_1^2}{t^2} = (\frac{dt^2 + dx_1^2}{t^2} - \frac{dt^2 + dx_1^2}{t^2 + \mathcal{O}(t^5)}) \rightarrow 0$ as $t \rightarrow 0^+$ which shows asymptotic convergence to g_H .

A similar analysis takes place for the solutions to (16) given by Lemma 5.1(ii), but this time $\rho(t) \rightarrow \infty$ as $t \rightarrow t_0^-$, showing the incompleteness of the metric. \square

REFERENCES

- [1] Baird, P., Wood, J.C., *Harmonic morphisms between Riemannian manifolds*, London Math. Soc. Monographs, New Series, vol. 29, Oxford Univ. Press, 2003.
- [2] Besse, A., *Einstein Manifolds*, Springer-Verlag, 1987.
- [3] Danielo, L., *Structures Conformes, Harmonicité et Métriques d'Einstein*, Ph.D. thesis, Université de Bretagne Occidentale, 2004.
- [4] Danielo, L., *Construction de métriques d'Einstein à partir de transformations biconformes*, Ann. Fac. Sci. Toulouse (6) **15** (3) (2006), 553–588.
- [5] Dieudonné, J., *Foundations of Modern Analysis*, Academic Press, 1969.
- [6] Hebey, E., *Scalar curvature type problems in Riemannian geometry*, Notes of a course given at the University of Rome 3. <http://www.u-cergy.fr/rech/pages/hebey/>.
- [7] Hebey, E., *Introduction à l'analyse non linéaire sur les variétés*, Diderot, Paris, 1997.
- [8] Hilbert, D., *Die Grundlagen der Physik*, Nachr. Ges. Wiss. Göttingen (1915), 395–407.
- [9] Hitchin, N.J., *On compact four-dimensional Einstein manifolds*, J. Differential Geom. **9** (1974), 435–442.
- [10] Schoen, R., *Conformal deformation of a Riemannian metric to constant scalar curvature*, J. Differential Geom. **20** (1984), 479–495.
- [11] Spivak, M., *A Comprehensive Introduction to Riemannian Geometry*, 2nd ed., Publish or Perish, Wilmington DE, 1979.
- [12] Thorpe, J.A., *Some remarks on the Gauss-Bonnet formula*, J. Math. Mech. **18** (1969), 779–786.
- [13] Vaisman, I., *Conformal foliations*, Kodai Math. J. **2** (1979), 26–37.
- [14] Yamabe, H., *On a deformation of Riemannian structures on compact manifolds*, Osaka Math. J. **12** (1960), 21–37.

LABORATOIRE DE MATHÉMATIQUES DE BRETAGNE ATLANTIQUE UMR 6205,
UNIVERSITÉ DE BREST,
6 AV. VICTOR LE GORGEU – CS 93837,
29238 BREST CEDEX, FRANCE
E-mail: paul.baird@univ-brest.fr

INSTITUTE OF MATHEMATICS,
UNIVERSITY OF THE PHILIPPINES DILIMAN,
C.P. GARCIA AVE.,
1101 QUEZON CITY, PHILIPPINES
E-mail: jventura@math.upd.edu.ph